

12-2018

An Investigation of Enzymes Capable of Activating Cannabinoid Prodrugs

Brittney J. Morgan

Follow this and additional works at: <https://digscholarship.unco.edu/theses>

Recommended Citation

Morgan, Brittney J., "An Investigation of Enzymes Capable of Activating Cannabinoid Prodrugs" (2018). *Master's Theses*. 69.
<https://digscholarship.unco.edu/theses/69>

This Text is brought to you for free and open access by the Student Research at Scholarship & Creative Works @ Digital UNC. It has been accepted for inclusion in Master's Theses by an authorized administrator of Scholarship & Creative Works @ Digital UNC. For more information, please contact Jane.Monson@unco.edu.

© 2018

BRITTNEY MORGAN

ALL RIGHTS RESERVED

UNIVERSITY OF NORTHERN COLORADO
Greeley, Colorado
The Graduate School

AN INVESTIGATION OF ENZYMES CAPABLE OF
ACTIVATING CANNABINOID PRODRUGS

A Thesis Submitted in Partial Fulfillment
of the Requirements for the Degree of
Master of Science

Brittney J. Morgan

College of Natural and Health Sciences
Department of Chemistry and Biochemistry

December 2018

This Thesis by: Brittney J. Morgan

Entitled: *An Investigation of Enzymes Capable of Activating Cannabinoid Prodrugs*

has been approved as meeting the requirements for the Degree of Master of Science in
the College of Natural and Health Sciences in the Department of Chemistry and
Biochemistry

Accepted by the Thesis Committee:

Richard M. Hyslop, Ph.D., Chair

Corina E. Brown, Ph.D., Committee Member

David L. Pringle, Ph.D., Committee Member

Accepted by the Graduate School

Linda L. Black, Ed.D.
Associate Provost and Dean
Graduate School and International Admissions

ABSTRACT

Morgan, Brittney J. *An Investigation of Enzymes Capable of Activating Cannabinoid Prodrugs*. Unpublished Master of Science thesis, University of Northern Colorado, 2018.

The plant *Cannabis sativa* contains a class of terpenophenols called cannabinoids which, in their active form, have been shown to have antiproliferative, anti-metastatic, anti-angiogenic, and pro-apoptotic effects on several types of cancer cells both *in vitro* and *in vivo*. Cannabinoids have been shown to not exert this apoptotic activity in healthy cells and they do not elicit the usual side effects seen with conventional chemotherapies. However, naturally occurring prodrugs from *Cannabis* must first be converted into the active form to induce the apoptotic and other physiological activities. Although humans do not possess the enzyme(s) necessary for this conversion, several different microbial species have been shown to have enzymes capable of catalyzing reactions similar to that required for activation of the cannabinoid prodrug(s).

This research focused on the isolation of the prodrug cannabidiolic acid (CBDA) from *Cannabis* and screening of various microbial cell lines capable of enzymatic conversion of CBDA to the active drug cannabidiol (CBD). This involved the development of a preparative HPLC method for isolation of CBDA and an analytical HPLC method for the monitoring of enzymatic prodrug conversion. Seven microbial species were screened for CBDA decarboxylation activity of the prodrug from *Cannabis*.

The cells were incubated with CBDA, and the substrate and resulting product(s) from incubation were simultaneously extracted from the incubated cell suspension at different time intervals and analyzed using HPLC. The Gram-negative bacterium *Pseudomonas putida* was identified to possess this enzymatic activity and will be an ideal species for gene isolation and further screening of the target enzyme. The results from this research provide an alternative to the traditional mode of activating cannabinoid pro-drugs and a new way to potentially control and monitor the effective dose of cannabinoids being administered.

ACKNOWLEDGEMENTS

First, I would like to thank my committee members. My research advisor Dr. Richard Hyslop for the many lessons learned throughout this project about my research, myself, and everything in between. In times of doubt you gave me confidence in myself and this project and ultimately, the will to succeed. Thank you to Dr. Corina Brown who gave me advice and guidance with my research but also had faith in my teaching abilities and fostered them with opportunities to teach lecture. My gratitude also extends to Dr. David Pringle, who provided a great deal of guidance with spectral analysis, pushed me to assess my work ethic with a gentle touch and took the time to be part of my thesis committee.

I would also like to thank Dr. Ann Hawkinson, Tyler Sherman, Doug Petty and our undergraduate collaborators in the School of Biological Sciences at UNC who assisted with the cell growth and harvest. Thank you to my fellow laboratory members Matt Brown, Shawn Bydalek, Sean Flora, Conner Hansen, Chris Laster, Anna Margiotta, Dr. Brown, and Dr. Hyslop for lending a hand in the laboratory in numerous ways, including assay preparation and instrumentation. A special thanks to Chad Wangeline and Casey Rogers for all their support with the instrumentation. Thank you to Pharmacyte Biotech for funding of this project. To my fellow graduate students, a heartfelt thank you for being there and supporting me through this experience; your comradery has made this experience rewarding and worthwhile. Finally, I want to thank

Mom, Dad, my partner Dana, and the rest of my family whose love and support have followed me from the very beginning of this adventure.

TABLE OF CONTENTS

CHAPTER

I.	INTRODUCTION.....	1
II.	REVIEW OF LITERATURE.....	8
	Cannabinoids	8
	Cannabinoid Physiology	12
	Cannabidiol and Cancer	16
	Enzymatic Decarboxylation	25
	Cell-in-a-Box	33
	Summary	37
III.	METHODOLOGY	39
	Chemicals and Reagents.....	39
	Cells	39
	Preparation of Solutions	40
	Construction of Calibration Curves	41
	Isolation of Cannabidiolic Acid from Cannabis Plant Material...	42
	Cell Culture	42
	Cell Assay for Decarboxylase Activity	43
	Instrumentation	44

IV.	RESULTS AND DISCUSSION.....	48
	Analytical Method for Monitoring the Conversion of Cannabidiolic Acid to Cannabidiol.....	48
	Isolation and Characterization of Cannabidiolic Acid.....	54
	Cell Assay for Decarboxylase Activity.....	69
V.	CONCLUSIONS	75
	Future Research.....	77
	REFERENCES.....	80
	APPENDIX	
A	CONDITIONS FOR INDIVIDUAL CELL ASSAYS FOR DECARBOXYLASE ACTIVITY.....	92
B	CHROMATOGRAMS OF FRACTIONS 2-6 COLLECTED FROM THE PREPARATORY HPLC METHOD.....	94
C	REPRESENTATIVE HPLC CHROMATOGRAMS FOR DECARBOXYLASE ACTIVITY IN VARIOUS CELL LINES	97
D	RIGHTS AND PERMISSIONS FOR USE OF FIGURES.....	127

LIST OF TABLES

TABLE

1	Area of Cannabidiolic Acid Peak at Different Concentrations Using the Analytical HPLC Method.....	52
2	Area of Cannabidiol Peak at Different Concentrations Using the Analytical HPLC Method.....	53
3	Summary of Collected Fractions from the Preparatory HPLC Method.....	59
4	Comparison of ¹ H- and ¹³ C-NMR Spectra of the Compound Isolated from Peak 1 to Assignments for Cannabidiolic Acid Reported by Choi et al. (2004).....	62
5	Percent Recovery Data.....	67
6	Summary of Cell Incubations for Decarboxylase Activity.....	70
7	Conditions for Individual Cell Assays for Decarboxylase Activity.....	93

LIST OF FIGURES

FIGURE		
1	Structures of Δ^9 -tetrahydrocannabinolic acid (THCA), cannabidiolic acid (CBDA), and cannabigerolic acid (CBGA)	2
2	Decarboxylation of cannabidiolic acid (CBDA) to cannabidiol (CBD).....	5
3	Biosynthesis of cannabichromenic acid (6), tetrahydrocannabinolic acid (7), and cannabidiolic acid (8) from the precursors IPP (1) and DMAPP (2).....	9
4	Representative structures of THC _s , CBD _s , CBN _s , CBG _s , CBC _s , and CBND _s	10
5	Numbering system for THC and CBD.....	11
6	Schematic of CB ₁ and CB ₂ cellular signaling.....	18
7	Schematic of various cell-death pathways potentially regulated by CBD.....	19
8	Cell survival (as measured by an MTT assay) of various breast cancer cell lines treated with increasing concentrations of CBD for 24 hours.....	23
9	Structures of Category 1 substrates.....	26
10	Structures of Category 2 substrates.....	28
11	Structures of Category 3 substrates.....	30
12	Structures of Category 4 substrates.....	31
13	Decarboxylation of benzoylformate to benzaldehyde.....	32
14	General schematic Cell-in-a-Box®/Ifosfamid technology.....	35

15	Chromatogram of 2.0 mg/mL CBDA standard using the analytical HPLC method.....	50
16	Chromatogram of 2.0 mg/mL CBD standard using the analytical HPLC method.....	50
17	Chromatogram of a mixture of CBDA and CBD using the analytical HPLC method.....	51
18	Standard calibration curve for cannabidiolic acid.....	52
19	Standard calibration curve for cannabidiol.....	53
20	Gradient Method 1.....	55
21	Chromatogram of CBDA extract from Soxhlet extraction using the preparatory HPLC method with Gradient 1.....	55
22	Gradient Method 2.....	56
23	Chromatogram of CBDA extract from Soxhlet extraction using the preparatory HPLC method with Gradient 2.....	56
24	Gradient Method 3.....	57
25	Chromatogram of CBDA extract from the Soxhlet extraction using the preparatory HPLC method with Gradient 3.....	57
26	Chromatogram of CBDA extract after two successive 1:2 dilutions with absolute ethanol from the Soxhlet extraction using Gradient 3.....	58
27	Chromatogram using the analytical HPLC method of the compound recovered from peak 1.....	60
28	CBDA standard (2.0 mg/mL) using the analytical HPLC method.....	60
29	Numbering scheme used by Choi et al. (2004) for CBDA...	61
30	¹³ C-NMR spectrum of the compound isolated from peak 1 in CDCl ₃	63
31	¹ H-NMR spectrum of the compound isolated from peak 1 in CDCl ₃	64

32	Deuterium exchange experiment for the compound isolated from peak 1.....	65
33	IR analysis of the compound isolated from peak 1.....	66
34	Chromatogram using the analytical HPLC method of the compound recovered from peak 7 using the preparatory HPLC method.....	67
35	Chromatogram of THCA standard (1.0 mg/mL)	68
36	Chromatogram generated using the analytical HPLC method of the cannabinoid mixture recovered from the 40-70 minute fraction using the preparatory HPLC method with Gradient 3.	68
37	Chromatogram of phytocannabinoid mixture standard using the analytical HPLC method.....	69
38	Chromatogram of CBDA after 0 hours of incubation with lysed <i>P. putida</i>	73
39	Chromatogram of CBDA after 1 hour of incubation with lysed <i>P. putida</i>	73
40	Chromatogram of CBDA after 2 hours of incubation with lysed <i>P. putida</i>	73
41	Chromatogram of CBDA after 3 hours of incubation with lysed <i>P. putida</i>	74
42	Chromatogram of CBDA after 4 hours of incubation with lysed <i>P. putida</i>	74
43	Chromatogram of the residue recovered representing peak 2 in the preparatory chromatogram.....	96
44	Chromatogram of the residue recovered representing peak 3 in the preparatory chromatogram	96
45	Chromatogram of the residue recovered representing peak 4 in the preparatory chromatogram	96
46	Chromatogram of the residue recovered representing peak 5 in the preparatory	97

47	Chromatogram of the residue recovered representing peak 6 in the preparatory chromatogram	97
48	Chromatogram after 0 hours of incubation of CBDA with lysed <i>B. subtilis</i> induced with 40 mM salicylic acid.....	99
49	Chromatogram after 1 hour of incubation of CBDA with lysed <i>B. subtilis</i> induced with 40 mM salicylic acid.....	99
50	Chromatogram after 2 hours of incubation of CBDA with lysed <i>B. subtilis</i> induced with 40 mM salicylic acid.....	100
51	Chromatogram after 3 hours of incubation of CBDA with lysed <i>B. subtilis</i> induced with 40 mM salicylic acid.....	100
52	Chromatogram after 4 hours of incubation of CBDA with lysed <i>B. subtilis</i> induced with 40 mM salicylic acid.....	100
53	Chromatogram after 0 hours of incubation of CBDA with lysed <i>B. subtilis</i> induced with an unknown concentration of CBDA.....	101
54	Chromatogram after 0.5 hour of incubation of CBDA with <i>B. subtilis</i> induced with an unknown concentration of CBDA.....	101
55	Chromatogram after 1 hour of incubation of CBDA with <i>B. subtilis</i> induced with an unknown concentration of CBDA.....	101
56	Chromatogram after 2 hours of incubation of CBDA with <i>B. subtilis</i> induced with an unknown concentration of CBDA.....	102
57	Chromatogram after 0 hours of incubation of CBDA with intact <i>E. aerogenes</i>	102
58	Chromatogram after 0.5 hour of incubation of CBDA with intact <i>E. aerogenes</i>	102
59	Chromatogram after 1 hour of incubation of CBDA with intact <i>E. aerogenes</i>	103
60	Chromatogram after 2 hours of incubation of CBDA with intact <i>E. aerogenes</i>	103
61	Chromatogram after 3 hours of incubation of CBDA with intact <i>E. aerogenes</i>	103

62	Chromatogram after 0 hours of incubation of CBDA with lysed <i>E. aerogenes</i>	104
63	Chromatogram after 0.5 hour of incubation of CBDA with lysed <i>E. aerogenes</i>	104
64	Chromatogram after 1 hour of incubation of CBDA with lysed <i>E. aerogenes</i>	104
65	Chromatogram after 2 hours of incubation of CBDA with lysed <i>E. aerogenes</i>	105
66	Chromatogram after 3 hours of incubation of CBDA with lysed <i>E. aerogenes</i>	105
67	CBDA standard for <i>K. pneumonia</i> assay.....	106
68	Chromatogram after 0 hours of incubation of CBDA with intact <i>K. pneumonia</i> induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.....	106
69	Chromatogram after 0.5 hour of incubation of CBDA with intact <i>K. pneumonia</i> induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.....	107
70	Chromatogram after 1 hour of incubation of CBDA with intact <i>K. pneumonia</i> induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.....	107
71	Chromatogram after 2 hours of incubation of CBDA with intact <i>K. pneumonia</i> induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.....	107
72	Chromatogram after 3 hours of incubation of CBDA with intact <i>K. pneumonia</i> induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.....	108
73	Chromatogram after 4 hours of incubation of CBDA with intact <i>K. pneumonia</i> induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.....	108
74	Chromatogram after 0 hours of incubation of CBDA with lysed <i>A. clavatus</i> induced with 4 mM salicylic acid.....	109
75	Chromatogram after 1 hour of incubation of CBDA with lysed <i>A. clavatus</i> induced with 4 mM salicylic acid.....	109

76	Chromatogram after 2 hours of incubation of CBDA with lysed <i>A. clavatus</i> induced with 4 mM salicylic acid.....	110
77	Chromatogram after 3 hours of incubation of CBDA with lysed <i>A. clavatus</i> induced with 4 mM salicylic acid.....	110
78	Chromatogram after 4 hours of incubation of CBDA with lysed <i>A. clavatus</i> induced with 4 mM salicylic acid.....	111
79	Chromatogram after 0 hours of incubation of CBDA with lysed <i>A. niger</i> induced with 4 mM salicylic acid.....	112
80	Chromatogram after 1 hour of incubation of CBDA with lysed <i>A. niger</i> induced with 4 mM salicylic acid.....	112
81	Chromatogram after 2 hours of incubation of CBDA with lysed <i>A. niger</i> induced with 4 mM salicylic acid.....	113
82	Chromatogram after 3 hours of incubation of CBDA with lysed <i>A. niger</i> induced with 4 mM salicylic acid.....	113
83	Chromatogram after 4 hours of incubation of CBDA with lysed <i>A. niger</i> induced with 4 mM salicylic acid.....	115
84	Chromatogram after 0 hours of incubation of CBDA with lysed <i>P. chrysogenum</i> induced with 4 mM salicylic acid.....	115
85	Chromatogram after 1 hour of incubation of CBDA with lysed <i>P. chrysogenum</i> induced with 4 mM salicylic acid.....	115
86	Chromatogram after 2 hours of incubation of CBDA with lysed <i>P. chrysogenum</i> induced with 4 mM salicylic acid.....	116
87	Chromatogram after 3 hours of incubation of CBDA with lysed <i>P. chrysogenum</i> induced with 4 mM salicylic acid.....	116
88	Chromatogram after 4 hours of incubation of CBDA with lysed <i>P. chrysogenum</i> induced with 4 mM salicylic acid.....	117
89	Chromatogram after 0 hours of incubation of CBDA with intact <i>P. putida</i> Induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.....	118

90	Chromatogram after 0.5 hours of incubation of CBDA with intact <i>P. putida</i> induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.....	118
91	Chromatogram after 1 hours of incubation of CBDA with intact <i>P. putida</i> induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.....	118
92	Chromatogram after 2 hours of incubation of CBDA with intact <i>P. putida</i> induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.....	119
93	Chromatogram after 3 hours of incubation of CBDA with intact <i>P. putida</i> induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.....	119
94	Chromatogram after 0 hours of incubation of CBDA with lysed <i>P. putida</i> induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.....	119
95	Chromatogram after 0.5 hours of incubation of CBDA with lysed <i>P. putida</i> induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.....	120
96	Chromatogram after 1 hour of incubation of CBDA with lysed <i>P. putida</i> induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.....	120
97	Chromatogram after 2 hours of incubation of CBDA with lysed <i>P. putida</i> induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.....	120
98	Chromatogram after 3 hours of incubation of CBDA with lysed <i>P. putida</i> induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.....	121
99	Chromatogram after 0 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	121
100	Chromatogram after 1 hour of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	122
101	Chromatogram after 2 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	122

102	Chromatogram after 3 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	122
103	Chromatogram after 4 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	123
104	Chromatogram after 5 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	123
105	Chromatogram after 6 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	123
106	Chromatogram after 0 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	124
107	Chromatogram after 1 hour of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	124
108	Chromatogram after 2 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	124
109	Chromatogram after 3 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	125
110	Chromatogram after 4 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	125
111	Chromatogram after 0 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	125
112	Chromatogram after 0.5 hour of incubation of CBDA with lysed <i>P. putida</i> induced 3 days with 10 mM salicylic acid.....	126
113	Chromatogram after 1 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	126
114	Chromatogram after 2 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	126
115	Chromatogram after 3 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	127
116	Chromatogram after 4 hours of incubation of CBDA with lysed <i>P. putida</i> induced with 10 mM salicylic acid.....	127

CHAPTER I

INTRODUCTION

With origins that date back to ancient times, many botanical extracts utilized as folk remedies have now been developed into modern day medicine. Acetylsalicylic acid (e.g., aspirin) was synthesized from a compound that came from the bark of Willow trees (*Salix alba*) which was used by many ancient cultures as a pain remedy before the discovery of salicylic acid as the naturally occurring active drug (Mahdi, Mahdi, Mahdi, & Bowen, 2006). More recently the plant *Cannabis sativa*, which has long been cultivated and harvested for its medicinal qualities, has been of interest to the scientific community. The first accounts of *Cannabis* used medicinally come from the world's oldest pharmacopoeia, the *Pên-tsao Ching*, which was compiled in 4,000 B.C. based upon oral traditions (Li, 1974). Despite its historic prevalence, however, the primary active compounds in the plant *Cannabis sativa*, phytocannabinoids, were not initially characterized until 1964 (Gaoni & Mechoulam, 1964).

At least 150 different phytocannabinoid compounds, considered terpenophenols, have been identified from *Cannabis sativa* over the past few decades (ElSohly & Gul, 2016). Of these compounds, the most abundant are Δ^9 -tetrahydrocannabinolic acid (THCA), cannabidiolic acid (CBDA), cannabigerolic acid (CBGA) (Figure 1), Δ^9 -tetrahydrocannabinol (THC), cannabidiol (CBD), cannabinol (CBN), and cannabigerol (CBG) (Figure 4) (Backer et al., 2009).




Figure 1: Structures of Δ^9 -tetrahydrocannabinolic acid (THCA), cannabidiolic acid (CBDA), and cannabigerolic acid (CBGA).

Initially, due to their lipophilic nature, it was assumed that cannabinoids exerted their effects by acting on biological membranes (Mavromoustakos, Yang, & Makriyannis, 1995); however, this hypothesis was challenged following the discovery of an endogenous cannabinoid system along with endogenous agonists (mainly anandamide and 2-arachidonoyl glycerol, termed endocannabinoids) of the cannabinoid receptors (Pacher, Bátkai, & Kunos, 2006).

The two major cannabinoid receptors in the endocannabinoid system have been classified as Cannabinoid Receptor Type 1 (CB₁) and Cannabinoid Receptor Type 2 (CB₂) (Pertwee, 2008a). The CB₁ receptor is more responsive to psychoactive cannabinoids, such as THC, than to non-psychoactive cannabinoids, such as CBD (Matsuda, Lolait, Brownstein, Young, & Bonner, 1990). Interestingly, there is a high concentration of CB₁ receptors in the brain. The crystal structure of the human CB₁


receptor was elucidated by Hua et al. in 2016, providing not only a further understanding of how the endogenous ligands bind but also a rationale for CB₁ drug design.

Although CB₂ receptors are found primarily in membranes of the immune system cells, such as those found in lymph tissues, they have been observed as being more highly expressed on various cancerous cells. Activation of G-coupled inhibitory proteins (G_i) associated with these cell membrane receptors leads to inhibition of adenylyl cyclase as well as many other signaling transduction pathways that modulate cell function (Alexander, Smith, & Rosengren, 2009). Cannabinoids, via these cannabinoid receptors, have the ability to regulate cellular signaling pathways which are critical for cell growth and survival. Therefore, cannabinoids may be useful in the treatment of cancer because of the high concentration of CB₂ receptors in tumor cell membranes (Alexander et al., 2009). With the emergence of more research on the endocannabinoid system and the role of cannabinoids as agonists in this system, the use of cannabinoids as therapeutic agents has become an attractive prospect.

Phytocannabinoids from *Cannabis sativa* have been reported to possess a variety of medicinal properties including antiproliferative activity in animal models toward specific solid tumors such as brain, pancreas, breast, prostate, colon, liver, and lung cancers (Massi et al., 2004). These antiproliferative properties are expressed when receptors of the endocannabinoid system are bound by specific phytocannabinoids. For example, CBD displays an unexpectedly high potency as an agonist of CB₁ and CB₂ in cells expressing these receptors; indeed, the manner which it interacts with CB₂ receptors provides a possible explanation for CBD's ability to interfere with immune cell chemotaxis and migration (Pertwee, 2008b).

Along with the ability of cannabinoids to inhibit tumor growth which has been observed *in vitro* and *in vivo* in several mice models, cannabinoids have been observed to demonstrate selective cellular toxicity, inducing cell death *in vitro* in human brain tumor cells at concentrations that are not harmful to normal cells (Salazar et al., 2009). This selective toxicity is significant because an effective anti-cancer treatment should induce apoptosis in tumor cells while not effecting healthy cells. Many cancer treatments in clinical use today do not possess this selective property and thus affect healthy cells producing unwanted side effects in contrast to cannabinoid-based treatments which potentially produce very few, if any, side effects (Robson, 2001). The potential for treatments with this selective nature with minimal side effects makes cannabinoids the interest in this study.

Currently, there are several commercial cannabinoid-based treatments available which utilize this cannabinoid activity in the body including Sativex®, Marinol®, and Epidiolex® (GW Pharmaceuticals, 2018). Sativex® is a THC-CBD-enriched, botanical extract for treating moderate to severe symptoms of multiple sclerosis including pain and spasm and was the first drug to have been approved by the FDA with two active ingredients in a single drug. Marinol®, a synthetic THC analog, alleviates nausea and emesis in cancer patients undergoing chemotherapy and has also been demonstrated to mitigate anorexia and cachexia associated with AIDS or chemotherapy. The most current drug approved by the FDA is the CBD-based drug Epidiolex® used to treat seizures associated with Dravet syndrome and tuberous sclerosis (GW Pharmaceuticals, 2018).



A simple way to decarboxylate these compounds is to heat them, usually accomplished by smoking or baking *Cannabis* plant material. However, this is not an ideal way to deliver a proper dose of the active drug to the site of the tumor. Active drugs comprise only a small portion of the *Cannabis sativa* plant; thus when absorbed through ingestion or inhalation most of the active drug will be distributed throughout the blood stream and only a small amount of therapeutic agent will actually reach the site of the tumor (Casarett & Doull, 2013). Additionally, if delivered in this manner one would need to consider the psychoactive properties of Δ^9 -THC, the concentration of which

varies from each strain of the plant, and how it would affect adult and child patients regarding cognitive function, reaction time, learning and memory functions.

Enzymatic decarboxylation of cannabinoids was explored in this study as an alternative method to heating. While humans do not possess the enzyme(s) necessary to catalyze the decarboxylation of cannabinoid compounds, several cell types, both prokaryotic and eukaryotic, have been reported to possess enzymes capable of catalyzing this specific reaction using substrates with functional groups similar to the cannabinoids (Grant & Patel, 1969; Santha, Rao, & Vaidyanathan, 1996; Kirimura, Gunji, Wakayama, Hattori, & Ishii, 2010; Snini et al, 2013). Many of these cell lines reported in the literature were explored in this study to identify an organism with the appropriate enzyme capable of this decarboxylase activity. Those organisms which exhibit the most potential will be considered for gene isolation of the target enzyme which could then be transfected into a eukaryotic cell of choice. The modified cell would then be encapsulated using Cell-in-a-Box® cellulose-based live cell encapsulation technology (PharmaCyte, 2017) and injected directly upstream from a tumor. This would allow for the appropriate dosage of a highly-targeted therapeutic drug to be converted directly to the active drug at the site of the tumor, making it an attractive cancer treatment.

To identify an enzyme that could activate the naturally occurring prodrug CBDA to its active drug form CBD, four different bacterial species and three fungal species were screened for activity; these species included *Enterobacter (Klebsiella) aerogenes*, *Pseudomonas putida*, *Klebsiella pneumonia*, *Bacillus subtilis*, *Aspergillus clavatus*, *Aspergillus niger*, and *Penicillium chrysogenum*. The microbial cells, both intact and lysed, were incubated with the cannabinoid prodrug. The cannabinoid substrate and

product(s) were simultaneously extracted from the incubated cell suspension at appropriate time intervals during incubation and analyzed using high-performance liquid chromatography (HPLC). The retention times of peaks in the chromatograms were compared to known standards to determine if the prodrug was activated by the species assessed.

In this study, identification and quantification of cannabidiol was assessed through the use of HPLC. The most common methods for qualitative and quantitative analysis of phytocannabinoids from *Cannabis sativa* include GC/FID and GC/MS (Klein, 2015); however, although these methods are rapid and well established, these methods are not useful for heat-labile compounds which can be decarboxylated via heat. Instead, an analytical HPLC method for monitoring conversion of CBDA to CBD and a preparatory HPLC method for isolation of CBDA from crude *Cannabis* extract were developed. This allowed for the initial steps of development of an enzyme-driven, cannabinoid-based drug to be performed in-house, from plant to active drug.

CHAPTER II

REVIEW OF LITERATURE

Cannabinoids can be classified into three major categories; endocannabinoids, phytocannabinoids, and synthetic cannabinoids (Pertwee, 2008a). Endogenous endocannabinoids are made in animals, including humans; in contrast, phytocannabinoids are plant-derived products from *Cannabis* while synthetic cannabinoids are laboratory produced. Although a brief mention of endocannabinoids is presented, this review focuses on the structure, function, and biological activity of phytocannabinoids. In addition, a broad review of many different *in vitro* and *in vivo* studies on various cancer cell lines is presented as a rationale for cannabidiol as a targeted cancer chemotherapeutic agent. Enzymatic conversion of cannabinoid model compounds is also discussed in support of an enzyme-based conversion of the inactive prodrug cannabidiolic acid to cannabidiol. Finally, Cell-in-a-Box® technology is discussed as a potential mechanism of delivery using a genetically-engineered cell containing the gene necessary for an enzyme capable of catalyzing the conversion of prodrug to active drug.

Cannabinoids

Phytocannabinoids from *Cannabis sativa* are synthesized in their acidic forms, from isopentyl pyrophosphate (IPP) and dimethylallyl pyrophosphate (DMAPP), the precursors of all terpenoids (Fellermeier, Eisenreich, Bacher, & Zenk, 2001). Although believed to be biosynthesized via the mevalonate pathway prior to the 1990s, it has been

suggested that the precursors IPP and DMAPP are derived predominantly via the deoxyxylulose phosphate pathway (Eisenreich, Rohdich, & Bacher, 2001). An overview of the biosynthesis of major cannabinoids from these precursors is shown in Figure 3. In general, IPP (1) and DMAPP (2) are condensed to form geranyl diphosphate (3) which then combines with olivetolic acid (4) to form cannabigerolic acid (5) from which the remaining cannabinoids are derived. Due to their origin from these two isoprene derivatives and their phenolic properties, cannabinoids can be categorized as terpenophenols.




Figure 3: Biosynthesis of cannabichromenic acid (6), tetrahydrocannabinolic acid (7), and cannabidiolic acid (8) from the precursors IPP (1) and DMAPP (2). The intermediates geranyl diphosphate (3), olivetolic acid (4), and cannabigerolic acid (5) are also shown. From Fellermeier et al. (2001), used with permission (Appendix C).

Currently, over 150 cannabinoids have been characterized from *Cannabis sativa* (ElSohly & Gul, 2016). These cannabinoids can be categorized based on structural similarities as tetrahydrocannabinols (THCs), cannabidiols (CBDs), cannabinols (CBNs), cannabigerols (CBGs), cannabichromenes (CBCs), and cannabinodiols (CBNDs), the most abundant of which are THCs, CBDs, and CBNs. Those which are deemed structurally unique cannabinoids are categorized as miscellaneous (ElSohly & Slade, 2005). Structures of representative compounds from each category are shown in Figure 4.




Figure 4: Representative structures of THCs, CBDs, CBNs, CBGs, CBCs, and CBNDs. From Thakur, Duclos, & Makriyannis (2005), used with permission (Appendix C).

Although there is much structural variance among these six major categories of cannabinoids, most are composed of 21 carbons which includes an n-pentyl chain on C-3 (THCs) or 5'-C (CBDs) of the aromatic ring. The variability of functional group

moieties accounts for the difference in physiological activity among cannabinoids (Thakur et al., 2005). This variability in structure also causes chemical nomenclature of certain cannabinoids to differ, for example the numbering of CBD compared to THC. These two structures are closely related, and CBD has been shown to cyclize into THC under acidic conditions (Gaoni & Mechoulam, 1964); however, due to their differences in functionality, they are numbered differently. Numbering for THC is therefore determined by a pyran ring; CBD relies on a terpene ring for numbering priority (Mechoulam & Hanuš, 2002). These numbering systems are given in Figure 5.




Figure 5: Numbering system for THC and CBD.

The remainder of this review will focus predominately on the related cannabinoids CBDA and CBD. CBD is distinguished by its aromatic and terpene rings, which lay nearly perpendicular to each other. Two independent forms, which differ mainly in the conformation of the pentyl side chain, were noted by Jones et al. (1977) when elucidating the crystal structure. The major torsion-angle difference is for the C (5')-C(1'')-C(2'')-C(3''), which is -57.2° and 179.2° for forms 1 and 2, respectively. Also, the benzene ring in the first form is planar and all substituents are coplanar while in the second form the aromatic ring is twisted slightly with a torsion angle of -4.1° and the 3'-OH bent slightly out of the plane of the ring. CBDA differs from CBD by a

carboxylic acid group on the 6' carbon *ortho* to the phenolic hydroxyl (Mechoulam & Hanuš, 2002). The decarboxylation of CBDA allows it to act as a prodrug where, upon decarboxylation, different biological effects can be noted (Ligresti, 2006).

Cannabinoid Physiology

The ability of cannabinoids to regulate the endocannabinoid system and the role of its receptors has been supported in numerous studies. Shortly after the characterizations of several phytocannabinoids, starting with the characterization of Δ^9 -THC by Raphael Mechoulam in the 1960s, researchers began to speculate how cannabinoids exert their effects in the body. Although it was originally assumed that cannabinoids exerted their effects by acting on biological membranes due to their lipophilic nature (Mavromoustakos et al., 1995), it was later discovered that the biological effects exerted by cannabinoids is facilitated through binding to endogenous cannabinoid-like receptors (Fonseca, Costa, Almada, Correia-Da-Silva, & Teixeira, 2013). This is not to say that cannabinoids only act upon the endogenous endocannabinoid system, merely that it is the primary mode of action.

Two major cannabinoid receptors of the endocannabinoid system have been identified as Cannabinoid Receptor Type 1 (CB₁) and Cannabinoid Receptor Type 2 (CB₂) (Pertwee, 2008a). The crystal structure of the human CB₁ receptor was elucidated by Hua et al. in 2016. The characterization of these receptors allowed for a more in-depth understating of ligand-receptor interactions and helps provide rationale for drug discovery. CB₁ receptors are highly expressed in nociceptive areas in the brain, the cerebellum, limbic system, and basal ganglia. This location of the CB₁ receptor makes it more responsive to psychoactive cannabinoids, such as THC, than to non-psychoactive

cannabinoids, such as CBD (Matsuda et al., 1990). Those receptors located in the brain are sparse in density in lower brainstem areas controlling cardiovascular and respiratory function, which may explain why high doses of cannabinoids are not lethal (Herkenham et al., 1990). CB₁ receptors have also been detected in spleen, eye, uterus, and testes (Guzmán, 2003).

In contrast, the CB₂ receptors are found mainly in peripheral tissues and tissues of the lymph nodes, tonsils, and spleen. Due to these locations, CB₂ is mainly a peripheral immunomodulation receptor with an important role in mediating pain, inflammation, and physiological defense (Pacher et al., 2006). The CB₂ receptor has also been observed in tumor cells. Both CB₁ and CB₂ receptors belong to the G-protein-coupled family of receptors and thus are mediated by activation of G_{i/o} alpha subunits (Turu & Hunyady, 2010). Activation of these G-coupled cell-membrane proteins may lead to inhibition of adenylyl cyclase as well as many other signaling transduction pathways that modulate cell function (Alexander, Smith, & Rosengren, 2009).

Shortly after the identification of the CB₁ and CB₂ receptors, research expanded from just studying cannabinoids in marijuana to looking for cannabinoid-like compounds produced in the body that act on the endogenous cannabinoid receptors. The endogenous agonists anandamide and 2-arachidonoyl glycerol were discovered, revealing that the primary physiological function of the endocannabinoid system was to bind these endogenous ligands (Pacher, Bátkai, & Kunos, 2006). These endogenous ligands (endocannabinoids) are made within the cell and are synthesized on-demand throughout the body from precursors in lipid membranes (Alger, 2013), so they can be used exactly when and where they are needed in the body, rather than being packaged and stored

similar to other signaling molecules. Endocannabinoids travel in a retrograde fashion to inhibit neurotransmitter release, making the main function in the brain to attenuate release of neurotransmitters (Guzmán, 2003). These receptors, along with the endocannabinoids themselves and their regulatory enzymes, make up the endocannabinoid system which is present in many chordates of the animal kingdom (Elphick, 2012). Activation of this system is associated with various processes including regulation of appetite, motor activity, memory, learning, emesis, and nociception (Fernández-Ruiz et al., 2007). More generally, it can be said that the role of the endocannabinoid system which has been observed is a role in homeostasis by controlling metabolic functions such as energy storage and transport and immune function (Pagotto, Marsicano, Cota, Lutz, & Pasquali, 2006).

In a 1998 report by Marzo et al., the functions of endocannabinoids were summarized as aids to “relax, eat, sleep, forget, and protect” with a focus on the endocannabinoids and their relation to the sensation of hunger. The amount of leptin, a hunger-inhibiting molecule, was shown to be inversely correlated with the number of endocannabinoids produced suggesting a relationship between the endocannabinoid system and appetite. If the endocannabinoid system is stimulated (by endo- or phytocannabinoids), an individual will have less leptin, and therefore will feel hungrier. These results interestingly align with anecdotal evidence from recreational *Cannabis* users who self-reported feeling hungrier when ingesting more and more *Cannabis*.

Unsurprisingly because of the large number of cannabinoid receptors in the brain, the endocannabinoid system is also reported to be involved in the regulation of memory (Alvares, Genro, Diehl, & Quillfeldt, 2008). A common effect self-reported by *Cannabis*

users is short-term memory impairment. In 2009, Mazzola et al. reported a significant impairment in the memory of rats when Δ^9 -THC was injected intra-peritoneally at concentrations of 3.0, 5.0, and 6.0 mg/kg 30 minutes prior to a passive-avoidance task. Long term potentiation in the hippocampus, a process essential for formation of storage of long-term memory, has been shown to be suppressed in mice treated with THC (Hampson & Deadwyler, 1999). The role of the endocannabinoid system in memory is also supported by the observation that CB₁ knockout mice show enhanced memory and long-term potentiation (Jiang, 2005). Although it is generally agreed upon that cannabinoids do have a regulatory role in memory, contradictory information on whether these effects impair or improve different phases of the memory processes exist simultaneously. Impairments in object recognition have been linked to dysregulation of the glutamatergic system (Nilsson, Hansson, Carlsson, & Carlsson, 2007) as well as hippocampal and para hippocampal dysfunction (Fernández-Ruiz et al., 2007). Interestingly, a study found that CBD interacts with the glutamatergic system (Hallak et al., 2011); CBD augmented the effects of an N-methyl-d-aspartate (NMDA) receptor antagonist in a human study, a receptor which is associated with memory and learning impairments. CBD was also shown to be protective against glutamate neurotoxicity (Hampson & Deadwyler, 1999). Finally, memantine, another NMDA receptor antagonist, improved object recognition in a transgenic Alzheimer Disease mouse model (Scholtzova et al., 2008) suggesting that CBD may improve recognition memory via the glutamatergic pathway. Although much has been learned about the endocannabinoid system, as of yet, it is still incompletely understood.

Cannabidiol and Cancer

This thesis research is grounded in much evidence from prior research into the effectiveness of cannabinoids, and more specifically cannabidiol, as cancer treatments. Cannabidiol is of specific interest as a cancer drug because it has many of the same antiproliferative effects as other cannabinoids without the psychotropic effects (Izzo, Borrelli, Capasso, Di Marzo, & Mechoulam, 2009). Cannabinoids could be useful in the treatment of cancer due to their ability to regulate cellular signaling pathways critical for cell growth and survival (Alexander et al., 2009). Expression of CB₁ and CB₂ receptors has been found in many types of cancer cells, although this does not necessarily correlate with the expression of these receptors in the tissue of origin (Guzmán et al., 2006). The fact that these receptors are upregulated in cancerous tissue and not in the surrounding tissue may explain the observation by McAllister et al. (2005) that cannabinoids can selectively inhibit proliferation and induce cancerous cell death at concentrations that do not harm normal cells.

The expression of CB₁ and CB₂ receptors in tumors of patients with hepatocellular carcinoma was studied by Xu et al. (2006). Patients were treated with embolization therapy to essentially starve the cancerous tumors of blood flow and essential nutrients. Patients then underwent a hepatectomy and liver samples were preserved for analysis. Their study concluded that when higher numbers of CB₁ and CB₂ receptors are expressed in cancerous tissue, there is a correlated improvement in prognosis in patients with hepatocellular carcinoma (Xu et al., 2006) and potentially many other cancerous cell lines. Therefore, cannabinoids may be useful in the treatment of cancer because of the high concentration of cannabinoid receptors in tumor cells (Alexander et al., 2009) which

are selectively targeted by cannabinoids. Studies on the effect of CBD on prostate, breast, glioma, and other cancer cell lines are reviewed later in this section.

In general, activation of CB₁ by cannabinoids minimizes membrane depolarization on presynaptic cells by inhibiting voltage-gated Ca²⁺ channels and activation of voltage-gated K⁺ channels (Turu & Hunyady, 2010). The result is a decrease in neurotransmitter release from presynaptic cells. Activation of the cannabinoid receptors in cancer cells has been reported to increase the levels of ceramide which has been shown to activate a mitogen-activated protein kinase (MAPK) pathway known as the extracellular-signal-regulated kinase (ERK) survival pathway (Ellert-Miklaszewska, Grajkowska, Gabrusiewicz, Kaminska, & Konarska, 2007; Ozaita, Puighermanal, & Maldonado, 2007). Activation of the ERK pathway usually results in cell proliferation, however continual stimulation causes arrest of the cell cycle and apoptosis. Cannabinoids have also been shown to exert anti-proliferative effects on cancer cells by causing an arrest of the cell cycle through the inhibition of adenylyl cyclase (Fonseca et al., 2013). This pathway can occur through activation of either CB₁ or CB₂ since they are both coupled to inhibitory G-proteins. Inhibition of adenylyl cyclase leads to lower concentrations of intracellular cAMP thus decreasing the activity of protein kinase A (PKA) and also activating the ERK pathway similar to the effect by increased levels of ceramide (Figure 6).




Figure 6: Schematic of CB₁ and CB₂ cellular signaling. From Katchan, David, and Shoenfeld (2016), used with permission (Appendix C).

Other potential pathways include transient receptor potential cation channel, subfamily V, member 1 (TRPV1) activation, downregulation of the vascular endothelial growth factor (VEGF) pathway, production of reactive oxygen species (ROS), induction of endoplasmic reticulum (ER) stress, and the activation of both intrinsic and extrinsic apoptotic pathways (Ramer, Merkord, Rohde, & Hinz, 2012). The mechanism of tumor cell inhibition produced by CBD was shown to depend on the type of cancer cell being analyzed (Massi, Solinas, Cinquina, & Parolaro, 2013). Importantly, of the cannabinoids CBD seemed to involve direct TRPV1 activation and/or CB₂ indirect activation (via fatty acid amide hydrolase), as well as induction of oxidative stress (Ligresti, 2006). Due to its many mechanisms of action, some of which are noted in Figure 7, cannabidiol is of interest in treating a variety of cancers. Activation of CB receptors can be responsible for

the increase in levels of ceramide or decrease of cAMP level via inhibition of adenylyl cyclase, both events lead to apoptosis mediated by caspase activities. Also, the ability for TRPV receptors activated by cannabinoids to cause an increase in the level of ROS and oxidative stress which leads to apoptosis has been observed. Although each mechanism may be activated differently, the process is highly interconnected and each event will lead to the same end effect, tumor cell death.




Figure 7: Schematic of various cell-death pathways potentially regulated by CBD. From Calvaruso, Pellerito, Notaro, & Giuliano (2012), used with permission (Appendix C)

Prostate cancer is the most common non-skin cancer among men in the United States (National Cancer Institute, 2017). The *in vitro* antitumor and anti-inflammatory activities of CBD in human prostate cancer cell lines were investigated by Sharma, Hudson, Adomat, Guns, & Cox (2014). Treatment of human prostate cancer cell lines

LNCaP and PC3 with *Cannabis* extracts containing 20–60 µg/mL CBD showed negligible activity at the lowest concentration (20 µg/mL) and massive induction of apoptosis at the highest concentration (60 µg/mL). This supports the need for a targeted approach to delivery and precision dosing when using CBD as a therapeutic drug.

Also examined in the Sharma et al. (2014) study was the effect of a *Cannabis* extract on androgen receptor transcripts. Because over-expression of androgen receptors in prostate cancer cells may promote cell growth, inhibiting this expression with the use of CBD may prove beneficial in preventing tumor growth. Treatment of LNCaP cells with *Cannabis* extract prepared in dimethyl sulfoxide (DMSO) containing 20, 40, 60, and 70 µg/mL CBD resulted in a marked decrease in androgen receptor expression when assessed at 24 hours post treatment. Although a significant decrease in expression was observed at all concentrations of CBD, the highest concentration produced the largest decrease in androgen receptor expression.

Additional studies of human prostate cancer cells *in vitro* and *in vivo* using mouse models found that the CB₂ receptor was involved in the inhibition of prostate cancer cell growth in both experimental models (Olea-Herrero, Vara, Malagarie-Cazenave, & Díaz-Laviada, 2009). Petrocellis et al. (2012) studied the effects of several pure cannabinoids (CBC, CBD, CBG, CBN, CBDA, CBGA, CBDV (cannabidivarin), CBGV (cannabigevarin), THC, THCA, THCV (Δ^9 -tetrahydrocannabivarin), THCA (Δ^9 -tetrahydrocannabivarin acid)) and those in *Cannabis* extracts on different human prostate carcinoma cell lines both *in vitro* and *in vivo*. Of the pure cannabinoids studied, CBD (1–10 µM) was the most effective inhibitor of tumor growth *in vitro*. *In vivo* studies were done with a *Cannabis* extract which contained 64.4% (w/w) CBD since it proved to be

even more potent than pure CBD alone when evaluating cell viability via an MTT assay in another portion of the study. Indeed, when the extract was administered to tumors generated in athymic mice from LNCaP cells, the growth of these tumors was reduced (Petrocellis et al., 2012).

Breast cancer is the most common cancer in women, regardless of race or ethnicity, in the United States (National Cancer Institute, 2017). Several cannabinoids were screened for their ability to reduce cell proliferation of different breast tumor cell lines (MCF7 and MDA-MB-231) *in vitro* and *in vivo* in a mouse model by Ligresti (2006). CBD exhibited the highest potency of cannabinoids studied, with IC_{50} values being $8.2 \pm 0.3 \mu M$ (MCF7) and $10.6 \pm 1.8 \mu M$ (MDA-MB-231). CBD selectively inhibited the growth of these two breast cancer cell lines *in vitro* at a range of concentrations (2.0-25 μM). Cannabidiol at a concentration similar to its IC_{50} did not affect the vitality of nontumor cell lines. Only at a concentration as high as 25 μM , which inhibited 100% of tumor cell growth, was viability of non-cancerous cells affected. Data regarding the extent to which the healthy cells were affected were not noted in the study. Mice with xenograft tumors obtained by subcutaneous injection of human MCF7 and MDA-MB-231 breast carcinoma cells were treated *in vivo* with either pure cannabidiol (5 mg/kg) or a cannabidiol-rich extract (6.5 mg/kg) and exhibited significantly smaller tumors in comparison with control mice.

In addition to confirming the *in vitro* observation by Ligresti (2006) of a reduction in proliferation of the human breast cancer cell line MDA-MB-231 when exposed to CBD, McAllister et al. (2007) reported a reduction in cancer cell proliferation in an additional breast cancer cell line, MDA-MB-436. Cells were treated with 0.1, 1.0 or 1.5

μM solutions of CBD for three days. Because the $1.5 \mu\text{M}$ solution was the most antiproliferative, the McAllister group examined the inhibitory effect of CBD at this concentration on Id-1 expression. Id-1, an inhibitor protein of basic helix-loop-helix transcription factors, has been shown to be a key regulator of the metastatic potential of breast and additional cancers (Benezra, Davis, Lockshon, Turner, & Weintraub, 1990). McAllister et al. (2005) previously demonstrated that reducing Id-1 led to significant reductions in breast cancer cell proliferation *in vitro* suggesting that reducing Id-1 expression could be a therapeutic strategy. Their 2007 study found that after a three-day treatment with CBD ($1.5 \mu\text{M}$, daily), the mRNA transcription of the Id-1 gene was significantly inhibited. Minn et al. (2005) had previously shown that CBD regulated the expression of key genes involved in the control of cell proliferation and invasion via the down-regulation of Id-1. This down-regulation may be one of the mechanisms by which CBD can reduce tumor aggressiveness. More recently, it was discovered that the Id-1 down-regulation was mediated by the up-regulation of extracellular-signal-related kinase (ERK) phosphorylation which produces the active isoform of the ERK protein (McAllister et al., 2012). Previously mentioned studies already showed that over stimulation of the ERK pathway contributed to arrest of the cell cycle and apoptosis, further supporting the inhibitory action of CBD through the ERK pathway.

An *in vitro* study which demonstrated the concentration-dependent nature of CBD as an effective breast cancer treatment was reported by Shrivastava, Kuzontkoski, Groopman, & Prasad (2011). These researchers found that CBD induced apoptosis in human breast cancer cell lines MDA-MB-231, MCF-7, SK-BR-3, and ZR-75-1 in a concentration-dependent manner (Figure 8). Moreover, it was found that at these

concentrations, CBD had little effect on MCF-10A cells, a non-tumorigenic human mammary cell line. These studies support the desirability of CBD as an anticancer agent by demonstrating a selective toxicity for breast cancer cells.




Figure 8: Cell survival (as measured by an MTT assay) of various breast cancer cell lines treated with increasing concentrations of CBD for 24 hours. From Shrivastava et al. (2011), used with permission (Appendix C).

Gliomas, or tumors of glial origin, are notably aggressive forms of cancer to which traditional forms of treatment such as radiation and chemotherapy are generally unresponsive (Philip-Ephraim, Eyong, Williams, & Ephraim, 2012). In a study by Massi et al. (2004), CBD (25 μM) was demonstrated to be effective in inhibiting U87-MG and U373 human glioma cell proliferation *in vitro* while not affecting healthy glial cells. *In vivo* treatment with CBD significantly reduced tumor growth of subcutaneously implanted U87 human glioma cells in mice when administered at a dose of 0.5 mg per mouse. In their study, it appeared that the addition of CBD to cell culture media during *in vitro* studies led to a dramatic drop of mitochondrial oxidative metabolism and

viability in the glioma cells. When adding the antioxidant α -tocopherol at a concentration of 10 μ M the antiproliferative effect of CBD was significantly prevented. Inhibitory effects were observed more rapidly at concentrations close to their IC_{50} values with effects from CBD (25 μ M) observed at 24 hours as compared to THC (1 μ M) observed within 2-5 days as reported by Sanchez et al. (1998)

In a study by Ciani et al. (2008), the cytotoxic effect of N-cyclopentyl-7-methyl-1-(2-morpholin-4-ylethyl)-1,8-naphthyridin-4(1H)-on-3-carboxamide (CB13), a synthetic CB_2 agonist, was investigated *in vitro* using the four colon cancer cell lines HCT8, SW480, HCA7, and HCT15 which had previously been reported to express the CB_2 receptor. The results of the study showed the growth of all cell lines was inhibited in a concentration-dependent manner. The CB_2 agonist also reduced growth of colon cancer cells *in vivo* in a mouse model (Ciani et al., 2008). Additionally, an extract of *Cannabis sativa* with a high content of CBD was demonstrated to reduce proliferation in colorectal cancer cells but not in healthy cells in a study by Romano et al. (2013). Proliferation was evaluated by an MTT assay in both healthy and cancerous cells at concentrations of CBD ranging from 0.3-5 μ M CBD for 24 hours. Twenty percent inhibition of tumor cell growth was reported with the highest concentration of CBD.

An *in vivo* study using athymic nude mice revealed a significant inhibition of A549 lung tumor metastasis in animals treated with a dose of 5 mg/kg CBD every 72 hours for 28 days as compared to vehicle-treated controls. The number of metastatic lesions in mice treated with CBD was one nodule after 28 days while the number of nodules in mice treated with only the vehicle was six, yielding an 84% inhibition of metastasis (Ramer et al., 2012).

Enzymatic Decarboxylation

The carboxylic acid group *ortho* to the hydroxyl group on the aromatic ring of CBDA must first be removed to elicit the desired physiological effects (Ligresti, 2006). Many studies have reported the expression of an enzyme capable of catalyzing the decarboxylation of phenolic carboxylic acids and other structurally similar acids by several eukaryotic and prokaryotic species which include many from the genera *Pseudomonas*, *Klebsiella*, *Bacillus*, *Aspergillus*, and *Penicillium*. In order to prioritize substrates based on similarity to CBDA, they were organized into four categories with Category 1 being the most similar and Category 4 being the least similar to the phenolic carboxylic acid moiety of CBDA. These compounds were of interest because of their similarity of the aromatic ring moiety, which interacts with the catalytic domain of these decarboxylase enzymes, to the aromatic moiety of CBDA. For proper enzyme-substrate binding to occur, there must be similarity between compounds so that important regions of the binding site can interact properly with the substrate in order for the desired reaction to occur.

Substrates such as salicylic acid, 6-methylsalicylic acid, 2,3-dihydroxybenzoic acid and gentisic acid (2,5-dihydroxybenzoic acid) have hydroxyl groups *ortho* to the carboxyl group, making them similar in structure to CBDA. Structures of these Category 1 substrates are presented in Figure 9. Cell lines capable of catalyzing decarboxylation of these Category 1 substrates are reported below.




Figure 9: Structures of Category 1 substrates.

Bacillus subtilis will express *padC* and *bsdBCD* genes when induced with salicylic acid (Duy et al., 2007). *PadC* encodes for a phenolic acid decarboxylase (PAD) while the *bsdBCD* operon encodes for phenylacrylic acid decarboxylase B and 4-hydroxybenzoate decarboxylase C and D (Duy et al., 2007). Salicylic acid, a Category 1 substrate, was unsurprisingly decarboxylated to phenol by the phenolic acid decarboxylase from *padC* but also was decarboxylated by the phenylacrylic acid decarboxylase encoded by *bsdBCD* even though salicylic acid is not, in fact, a phenylacrylic acid (Duy et al., 2007). However, it is known that genes for these enzymes are down-regulated when stress is absent (Duy et al., 2007) and so induction was necessary.

Trichosporon moniliform was shown to produce the enzyme salicylic acid decarboxylase, which will catalyze the decarboxylation of not only salicylic acid but

other carboxylic acids as well (Kirimura et al., 2010). The enzyme also catalyzed the decarboxylation of b-resorcylic acid (2,4-dihydroxybenzoic acid) producing resorcinol (1,3-dihydroxybenzene), c-resorcylic acid (2,6-dihydroxybenzoic acid) producing resorcinol, 2,3-dihydroxybenzoic acid producing catechol (1,2-dihydroxybenzene), and 4-aminosalicylic acid producing 3-aminophenol.

Multiple fungal representatives have been reported to express a variety of decarboxylases. A well-classified enzyme transiently expressed in *Aspergillus* recognizes 6-methylsalicylic, a substrate containing a carboxylic acid *ortho* to a phenolic hydroxyl and a carbon side chain on C-6 which would be analogous to the pentyl chain on C-5' of CBD. The gene *PatG*, encoding this enzyme, is expressed by many fungi including *Aspergillus clavatus*, *Aspergillus giganteus*, and *Aspergillus longivesica* as well as *Byssoschlamys nivea*, and *Penicillium expansum* (Snini et al., 2013).

The non-oxidative decarboxylation of 2,3-dihydroxybenzoic acid (2,3-dHBA) to catechol is catalyzed by the enzyme 2,3-dihydroxybenzoic acid decarboxylase (2,3-dHBAD). This enzyme is produced by several *Aspergillus* fungi including *Aspergillus oryzae* and *Aspergillus niger*. Stable at physiological pH, dHBAD from *Aspergillus* shows high specificity towards 2,3-dHBA and does not decarboxylate its analog 2,3,4-trihydroxybenzoic acid (2,3,5-THBA). Interestingly, dHBAD from *Trichosporon cutaneum* exhibited a broad substrate specificity and decarboxylated 2,3,5-THBA unlike dHBAD from the *Aspergillus* species (Santha et al., 1996).

Halvorson (1963) demonstrated that a species of *Aspergillus* successfully decarboxylated 2,4-dihydroxybenzoic acid to form resorcinol as an intermediary step in

the metabolism of 2,4-dihydroxybenzaldehyde. The 2,4-dHBA was metabolized to form resorcinol after 17 days and was completely metabolized after 29 days.

Category 2 substrates such as 3,4,5-trihydroxybenzoic acid (gallic acid), 3,4-dihydroxybenzoic acid (o-protocatechuic acid), and 4-hydroxybenzoic acid (p-hydroxybenzoic acid) (Figure 10) have a phenolic hydroxyl *para* to the carboxylic acid group but lack the *ortho* hydroxyl group. Decarboxylases for these Category 2 substrates are still of interest, however, due to the variability of enzyme specificity.




Figure 10: Structures of Category 2 substrates.

An enzyme able to decarboxylate *o*-pyrocatechuic acid (3,4-dihydroxybenzoic acid), a Category 2 substrate, was purified from a member of the *Aspergillus* genus, *Aspergillus niger* (Subba, Moore, & Towers, 1967). *Klebsiella aerogenes*, previously known as *Enterobacter aerogenes*, expresses *p*-hydroxybenzoic acid decarboxylase (PHBD) which decarboxylates gentisic acid (Category 1), *p*-hydroxybenzoic acid (PHB), gallic acid (3,4,5-trihydroxybenzoic acid), and protocatechuic acid. Two days were required to decarboxylate a majority of PHB, gallic acid, and protocatechuic acid and four days for gentisic acid. A permeability barrier was observed for PHB due to the

increase in activity between intact cells and those which had been lysed with sonication. However, the optimum conditions for this enzyme (pH 6) was below human physiological pH (Grant & Patel, 1969).

Nakajima, Otani, & Niimura (1992) isolated *K. pneumoniae* and *Streptococcus faecalis* for the study of decarboxylase enzymes. In their study, they were able to demonstrate that, at an optimal pH of 6, *K. pneumoniae* possessed an enzyme capable of catalyzing the decarboxylation of gallic acid, a Category 2 substrate. Similarly, a decarboxylase enzyme from *S. faecalis* was able to decarboxylate both gallic acid and protocatechuic acid. Optimal conditions for this decarboxylase were found to be a pH of 6.6 and a temperature of 45 °C. Researchers were able to show that for every one molecule of gallic acid, one molecule of 1,2,3-trihydroxybenzene was formed, suggesting a 1 to1 relationship between the substrate gallic acid and its decarboxylated product from the decarboxylase enzyme partially purified from *S. faecalis*.

Three strains of *Lactobacillus brevis* were shown to decarboxylate gallic and protocatechuic acid (Curiel, Rodríguez, Landete, Rivas, & Muñoz, 2010). Cultures were incubated with a variety of 1 mM phenolic acids for 10 days at 30 °C to induce the decarboxylase. Only those incubated in the presence of gallic acid decarboxylated gallic acid and those incubated in the presence of protocatechuic acid decarboxylated protocatechuic acid. Following the observation of decarboxylase activity from *L. brevis*, a phenolic acid decarboxylase (*PAD*) gene was identified, cloned and characterized by Landete et al. (2010) showing an optimum activity at pH 6 and 22 °C.

Compounds which have well-characterized decarboxylases but are lacking much of the structural similarity to CBDA as in Category 1 and 2 are classified as Category 3

substrates such as *p*-coumaric acid, ferulic acid, and caffeic acid (Figure 11), all of which are derivatives of cinnamic acid.

Decarboxylases encoded by *PadC* in *Bacillus subtilis* were shown to display substrate-specificity for decarboxylation of ferulic, *p*-coumaric, and caffeic acid. These three acids are the three substrates of PAD (Duy et al., 2007). Studies from Cavin, Dartois and Divies (1998) showed that induction of the PAD enzyme must occur for decarboxylase activity to be observed in *B. subtilis*. Cell cultures were induced with ferulic, *p*-coumaric, and caffeic acid. All substrates were shown to induce decarboxylase activity, with caffeic acid having the lowest activity.




Figure 11: Structures of Category 3 substrates.

Category 4 encompasses miscellaneous substrates which have the least structural similarities to CBDA. A mixed culture of marine bacteria isolated from marine sediment (ON-7), *Bacillus subtilis* (Strain FO) and *Pseudomonas testosterone* were all cultured to examine the metabolism of phthalic acids (Taylor & Ribbons, 1983). The mixed culture

ON-7 was incubated with various phthalic acids for 4 hours at 30 °C. The culture was able to decarboxylate phthalic acid, 4-hydroxyphthalic acid, 3-fluorophthalic acid, 4-chlorophthalic acid, and 4,5-dihydroxyphthalic acid (Figure 12). *B. subtilis* was also incubated with various phthalic acids for 20 hours at 30 °C and was able to decarboxylate all substrates except the 4,5-dihydroxyphthalic acid. The *P. testosterone* culture was also incubated with phthalic acids for 24 hours at 30 °C; however, this cell line only decarboxylated 4-hydroxyphthalic acid and 4,5-dihydroxyphthalic acid.




Figure 12: Structures of Category 4 substrates.

An enzyme which has been shown to remove the carboxylic acid *ortho* to a hydroxyl group from 1,2-dihydro-1,2-dihydroxybenzoic acid (Figure 12) is 1,2-dihydro-1,2-dihydroxybenzoic acid dehydrogenase (Reiner, 1971). This enzyme catalyzes the final step in the conversion of 1,2-dihydro-1,2-dihydroxybenzoic acid to catechol and has been characterized from *Pseudomonas putida*, *Pseudomonas cepacia*, *Acinetobacter calcoaceticus*, and *Alcaligenes eutrophus* (Reiner, 1971).

Another enzyme which has been characterized from *P. putida* in addition to that for 1,2-dihydro-1,2-dihydroxybenzoic acid is benzoylformate decarboxylase (BFD) which catalyzes the decarboxylation of benzoylformate to benzaldehyde (Figure 13) (Iding et al., 2000). Although seeming highly specific to the substrate benzoyl formate, the decarboxylase showed minimal activity towards other carboxylic acid groups, in a pH range viable for the human body (pH 7). Important to note is that the purified enzyme did show stability at temperatures up to 60 °C for 2 hours.




Figure 13: Decarboxylation of benzoylformate to benzaldehyde.

In addition to the aforementioned body of literature, research conducted in our laboratory supports the decarboxylation of cannabinoid model compounds (Cribbs, 2014). Three bacterial species, *K. pneumonia*, *E. aerogenes*, and *P. putida* displayed decarboxylase activity for various substrates. *P. putida* was incubated with 2-hydroxy-4-methoxybenzoic acid for one hour, after which time the presence of the decarboxylation product, 3-methoxyphenol, was observed with intact cells. In contrast, incubation of *E. aerogenes* in the presence of PHB produced the decarboxylation product, phenol, after 20 minutes with both intact and lysed cells. In this case, induction was not necessary prior to incubation for decarboxylase activity; however, the amount of phenol produced was less than when induced. Finally, *K. pneumonia* was observed to decarboxylate PHB to

phenol following incubation for 30 minutes. After 60 minutes a marked decrease in PHB and increase in phenol was shown (Cribbs, 2014).

Based on the reviewed literature and previous work by our research group, seven cell lines were chosen for screening: *Aspergillus niger*, *Aspergillus clavatus*, *Bacillus subtilis*, *Enterobacter (Klebsiella) aerogenes*, *Klebsiella pneumonia*, *Penicillium chrysogenum*, and *Pseudomonas putida*. *K. pneumonia*, *E. aerogenes*, and *P. putida* displayed decarboxylase activity for various model compounds in previous studies by our research group and thus were carried forward to be screened with actual cannabinoids. *Bacillus subtilis* has a well-characterized PAD enzyme which decarboxylates a variety of compounds from Categories 1-4 and so was also chosen to be screened with cannabinoids. The same rationale was used in choosing those cell lines from the *Aspergillus* and *Penicillium* genera. The cells chosen contained decarboxylases which covered a variety of structural variations from that of CBDA to have a more complete understanding of which enzyme binding sites might be most compatible with the structure of CBDA.

Cell-in-a-Box®

The Cell-in-a-Box® live cell encapsulation technology was developed by Austranova, an Austrian biotechnology company, and employed by PharmaCyte Biotech as a delivery system of cells which will potentially convert prodrugs to active drugs for the treatment of a variety of different diseases. This delivery system is currently in clinical trials for treatment of pancreatic and breast cancer, as well as diabetes. PharmaCyte Biotech's model for the treatment of cancer involves encapsulating genetically-engineered human cells that convert an inactive chemotherapy drug into its

active or “cancer-killing” form (PharmaCyte, 2017). Cells are encapsulated through a reaction of two polymer coating steps in which the cells are mixed with one polymer and then the mixture is introduced to the next polymer using a droplet-forming machine, where the capsule membrane complex forms immediately around the droplets containing cells, resulting in capsules with a diameter of 0.7-0.8 mm which can encapsulate as many as 10,000 live cells (PharmaCyte, 2017).

The small, spherical capsules, made of a bio-inert cellulose-based polymer, are designed to retain the engineered cells due to size, while preventing immunoglobulins from entering the capsule and destroying the cells. The porous nature of the capsule allows blood flow, which nourishes the living cells inside, and does not block circulation when inserted into the blood stream. In cancer therapies, capsules are implanted as close as possible to the patient’s cancerous tumor which can be done so safely by using needles or catheters. Capsules that have been placed in the body for over two years have not degraded nor caused damage or inflammation to the surrounding tissues (PharmaCyte, 2017).

Cell-in-a-Box® technology is currently being implemented in clinical trials for the treatment of pancreatic cancer with the prodrug ifosfamide, a highly effective chemotherapy drug. However, ifosfamide is typically given intravenously and activated in the liver by cytochrome P450; consequently, the effective dose needed for the desired anti-cancer effects of ifosfamide can cause serious side effects. With the use of Cell-in-a-Box® technology, ifosfamide is given intravenously and converted to its active form when it comes into contact with the encapsulated cells implanted in the blood supply near the tumor instead of being converted in the liver and distributed throughout the body until

reaching the tumor site. A pictorial schematic of this mechanism of action is presented in Figure 14. In clinical trials, patients were able to receive a dose two-thirds less than the usual, which decreased or completely eliminated the undesirable side effects.




Figure 14: General schematic of Cell-in-a-Box®/Ifosfamid technology.

In a phase 1 and 2 clinical trial, 14 patients were treated with the Cell-in-a-Box®/Ifosfamide therapy. Cell-in-a-Box® capsules, with engineered cells which expressed the P450 system like that of the liver where ifosfamid is usually metabolized, were implanted on day zero. One dose of ifosfamide was given intravenously to patients during the first week (days 2-5) and a second dose was given during the fourth week (days 25-27), both at a dose of 1 g/m². Individual results varied from 25-50% reduction in tumor size; however, no patients showed tumor growth and two patients' inoperable tumors (due to size and location in the pancreas) became operable. These results were compared with phase 3 historical data for gemcitabine, a pancreatic cancer treatment currently approved by the FDA. The percentage of one-year survivors using the Cell-in-a-Box® therapy was double that previously reported for patients using gemcitabine, the median survival of patients was substantially increased as compared to gemcitabine, and

in contrast to gemcitabine, no treatment-related side effects were seen with Cell-in-a-Box®/Ifosfamid treatment (Pharmacyte, 2017).

In a phase 2 clinical trial using the Cell-in-a-Box®/Ifosfamid treatment, 13 pancreatic cancer patients were administered Cell-in-a-Box® technology in the same manner as in the phase 1/2 trials, however a dose of 2 g/m² was given. The conclusion was that a dose of 1 g/m² was more appropriate, since that dose was able to effectively treat the pancreatic tumors while still minimizing negative effects of the typical mode of delivery for ifosfamid. A future trial, awaiting FDA approval, is designed to show that Cell-in-a-Box® plus low-dose ifosfamide therapy can serve as an effective and safe consolidation chemotherapy for patients whose tumors no longer respond after four to six months of therapy with Abraxane® plus gemcitabine (PharmaCyte, 2017).

In addition to trials with human pancreatic cancer, a veterinary phase 1/2 trial using dogs with breast cancer has been conducted. Cell-in-a-Box® capsules containing cells with high CYP2B1 activity were used in combination with cyclophosphamide. Because it is often used to treat breast cancer in humans as well as mammary tumors in dogs, cyclophosphamide was chosen rather than ifosfamide. As in the pancreatic cancer clinical trials in humans, the capsules showed no side effects or irritation at the implanted sites (PharmaCyte, 2017).

A total of 16 dogs were enrolled in the preclinical trial. Ten dogs were treated with the Cell-in-a-Box®/cyclophosphamide combination, one of which had two tumors. Six of the 11 tumors from the dogs treated with the encapsulated cells/cyclophosphamide combination showed a partial remission (PR, regression of initial tumor mass by less than 50%) while the other five tumors showed stable disease (SD, regression of the initial

tumor mass by less than 50% or its enlargement by less than 25%). One group of six dogs, one of which had two tumors, was treated intravenously with cyclophosphamide alone at a dose of 7 mg/kg of body weight. Of the seven tumors in the six dogs that received cyclophosphamide alone, six tumors showed SD and the other exhibited progression of disease (PD, increase in initial tumor mass of at least 25% or the occurrence of new tumors). The median tumor size reduction of the tumors in the ten dogs treated with the combination Cell-in-a-Box®/cyclophosphamide was 53%. This is in contrast with only a 21% reduction of the size of the tumors in the dogs treated with cyclophosphamide alone (Michalowska et al., 2014). The results of this study suggest that the combination of the Cell-in-a-Box®/cyclophosphamide treatment may prove to be beneficial not only for the treatment of mammary cancers in the veterinary setting but may also be translated into an effective treatment of breast cancer in humans. This supports the application of the Cell-in-a-Box® technology as a treatment platform.

Summary

With the support from the studies reviewed in this chapter, this research aimed to isolate the phytocannabinoid cannabidiolic acid from *Cannabis sativa* plant material and to screen microbial cell lines which could potentially decarboxylate CBDA to CBD. With the broad review of many different *in vitro* and *in vivo* studies citing the effects of CBD on various cancer cell lines, this natural product clearly could be the basis of a targeted cancer therapy. Cannabidiol is already an FDA-approved drug in other medicinal contexts and its high potential for other therapies could be harnessed if an appropriate manner of administration was developed. One approach involving Cell-in-a-Box® begins with identifying a microbial cell line capable of enzymatic conversion of

the inactive prodrug cannabidiolic acid to cannabidiol by performing cell assays on a variety of species from the literature discussed. Ultimately, Cell-in-a-Box® technology could provide a potential mechanism of delivery for a genetically-engineered cell containing the gene necessary for an enzyme capable of catalyzing the conversion of prodrug to active drug.

CHAPTER III

METHODOLOGY

Chemicals and Reagents

Cannabidiolic acid (CBDA) (1244-58-2), cannabidiol (CBD) (13956-29-1), and phytocannabinoid mixture 1 (18791) were obtained from Cayman Chemical Company (Ann Arbor, Michigan). Chloroform-d (865-49-6) and formic acid (64-18-6) were obtained from Acros Organics (Geel, Belgium). HPLC-grade acetonitrile (75-05-8), absolute ethanol (64-17-5), n-pentane (109-66-0) and potassium hydroxide (1310-58-3) were obtained from Fisher Scientific (Fair Lawn, New Jersey). Ammonium formate (540-69-2), HPLC-grade methanol (67-56-1), Δ^9 -THC (1972-08-3) and salicylic acid (69-72-7) were obtained from Sigma-Aldrich Chemical Company (St. Louis, MO). The sources for the remaining chemicals and reagents are as follows: Tryptic soy broth (470015-844) (BD Sparks, MD); sabouraud-dextrose broth (1466366) (EMD, Darmstadt, Germany). *Cannabis* plant material (RTI log number 13784-1114-18-6, reference number SAF 027355, 6.5% CBD(A), 3.5% THC(A)) was obtained from the National Institute on Drug Abuse. HPLC-grade water (18.2 M Ω) was purified in-house by passing reverse osmosis purified water through a Milli-Q® Advantage A 10 Ultrapure Water Purification System.

Cells

Seven cell lines were screened for decarboxylase activity with CBDA.

Aspergillus clavatis (1007) and *Apergillus niger* (16888) were purchased from the

American Type Culture Collection (Manassas, VA). *Bacillus subtilis* (470176-524), *Enterobacter (Klebsiella) aerogenes* (470179-294), *Klebsiella pneumoniae* (470179-122), *Penicillium chrysogenum* (470176-388) and *Pseudomonas putida* (470179-092) were obtained from VWR International (Randor, PA) via Ward's Science Co. (Rochester, NY).

Preparation of Solutions

All mass measurements were made using an Ohaus Explorer Pro analytical balance. The pH of buffer solutions was measured using a Fisher Scientific Accumet[®] AB150 pH meter. The meter was calibrated using buffers obtained from Fisher Scientific at pH 4.0 (SB-101-500), 7.0 (SB-107-500), and 10.0 (SB-115-500). All solvents were filtered prior to use with the HPLC. Aqueous solvents and methanol were filtered through a nylon filter (0.22 μ m) and all other organic solvents were filtered through a Teflon filter (0.22 μ m).

Ammonium formate (20.0 mM, pH 3.20) preparation. To prepare 1.00 L of 20.0 mM ammonium formate, 1.2612 g of ammonium formate were added to a 1000-mL volumetric flask and diluting to volume with HPLC-grade water. Formic acid (20.0 mM) was prepared by adding 0.3835 mL of 96% formic acid to a 500-mL volumetric flask and diluted to volume with HPLC-grade water. The 20.0 mM formic acid solution was added to the 20 mM ammonium formate solution until a pH of 3.20 was observed on the pH meter.

Cannabidiolic acid (2.0 mg/mL) stock solution preparation. To prepare 700 μ L, 1.4 mg of CBDA were added to a vial and dissolved in 700 μ L of absolute ethanol, added by a micropipette.

Cannabidiol (2.0 mg/mL) stock solution preparation. To prepare 1.05 mL, 2.1 mg of CBD were added to a vial and dissolved in 1.05 mL absolute ethanol, added by micropipette.

Standard curve solutions for cannabidiolic acid. Five 200 μ L standards with concentrations of 0.10, 0.50, 1.0, 1.5, and 2.0 mg/mL were prepared by making appropriate dilutions of the stock solution using a variety of Rainin Pipetman micropipettes. The standards were diluted to a volume of 200 μ L with absolute ethanol. All standards were prepared in triplicate.

Standard curve solutions for cannabidiol. Five 200 μ L standards with concentrations of 0.10, 0.50, 1.0, 1.5, and 2.0 mg/mL were prepared by making appropriate dilutions of the stock solution using a Rainin Pipetman micropipette. The standards were diluted to a volume of 200 μ L with absolute ethanol. All standards were prepared in triplicate.

Induction solution preparation. The appropriate mass of the induction substrate, according to assay, was dissolved in an appropriate solvent in a volumetric flask depending on the substrate and the desired volume. For example, salicylic acid was prepared as the potassium salt by dissolving the appropriate amount in an equimolar + 10% equivalent of KOH in a 250-mL volumetric flask. An additional 25 mL of absolute ethanol were added then diluted to volume with HPLC-grade water.

Construction of Calibration Curves

Calibration curves for both CBDA and CBD were constructed for use to determine the concentration of both CBDA present and CBD formed, if any, from the signal produced during HPLC analysis. A 2.0 mg/mL stock solution of each compound

was prepared and then diluted with absolute ethanol to make solutions of 1.5, 1.0, 0.50, and 0.10 mg/mL. Each standard was prepared in triplicate and analyzed using the analytical HPLC method. The area under the curve of the analyte peak was averaged across the triplicate solutions and plotted in Microsoft Excel against concentrations to generate a standard curve.

Isolation of Cannabidiolic Acid from Cannabis Plant Material

Initially, 5.00 g of *Cannabis* plant material, containing a reported 6.5% CBD(A) content, obtained from the National Institute for Drug Abuse was freeze-dried and pulverized with a mortar and pestle prior to extraction. The dry material was placed in a filter paper thimble which was then placed into the main chamber of a Soxhlet extractor. The Soxhlet extractor was placed onto a round-bottom flask containing 100 mL of pentane which was then heated to a gentle reflux in a continuous solid-liquid extraction for three hours. After extraction, the pentane was removed by rotary evaporation to recover a dark green-brown, oily and viscous extract containing CBDA and other cannabinoids. The resulting oil was resuspended in 15 mL ethanol and gravity filtered to remove any insoluble solids. CBDA was isolated from the resulting solution as described in the Preparative Method for Purifying CBDA from a Crude *Cannabis* Extract outlined in the Instrumentation section of Methodology.

Cell Culture

Cells were cultured in 500-700 mL of appropriate growth medium and incubated at room temperature or 37 °C according to species. Cells were induced for 24-72 hours with model compounds known for inducing decarboxylase activity. Treatments for each assay are listed in Table 9 (Appendix A). For example, salicylic acid was added to a

culture of *Pseudomonas putida* to give a final concentration of 1 mM and incubated for 72 hours to induce expression of decarboxylase enzymes. Following induction, cells were centrifuged in 50-mL conical tubes at 4000g for 10 minutes to collect all cells in a pellet. Wet cell mass of the final cell pellet was obtained in mg/mL. Cells were resuspended in 30 mL of fresh culture medium and divided between two 50-mL conical tubes. One sample was sonicated to lyse the cells or used intact for assessing decarboxylase activity. Sonication was performed using a Branson Digital sonifier using a program designed to sonicate the sample at 40% amplification using one second bursts followed by a one second pause for a total of twenty seconds. Cells were on ice during sonication to ensure that the sample remained cool. The other sample, used as control, was autoclaved. All samples were kept on ice until screened for enzymatic activity.

Cell Assay for Decarboxylase Activity

Each strain of microbial cells was grown according to the procedure outlined in the Cell Culture section of Methodology. Cells were assessed for enzymatic decarboxylation of CBDA alongside an autoclaved control. Cells were assayed at a range of concentrations of CBDA from 0.23-0.36 mM by making solutions of 1.1-1.75 mg of CBDA in 100 μ L of absolute ethanol and adding to the incubation medium. Specific concentrations of CBDA are given in Table 9 (Appendix A). Addition of the CBDA solution (100 μ L) to the incubation flask was followed by the addition of 15.0 mL of cell suspension. The solutions were incubated in a constant-temperature, shaking water bath from VWR International at 36 °C and shaken at 60 rpm for the duration of the assay. Incubation media was removed in 2.0 mL aliquots at time = 0, 1, 2, 3, 4, 5, and 6 hours and immediately extracted with 5.0 mL of pentane containing 0.5 mL absolute ethanol by

vortexing with the 2.0 mL aliquot of incubated media. The cannabinoid-containing pentane layer was separated from the cell suspension using centrifugation; the pentane layer was transferred into a separate vial and evaporated to dryness using nitrogen gas. The residue was resuspended in 100 μ L of absolute ethanol. A 20 μ L aliquot of the ethanol solution was loaded into the HPLC 10 μ L loop injector and analyzed for the presence of CBD using HPLC in accordance to the Analytical Method for Monitoring Conversion of CBDA to CBD outlined in the Instrumentation section of Methodology.

Instrumentation

Analytical High Performance Liquid Chromatographic Method for Monitoring Conversion of Cannabidiolic Acid to Cannabidiol

HPLC analyses were performed using a Shimadzu (Tokyo, Japan) VP equipped with a SPD-M10A diode array detector, a Shimadzu SCL-10A system controller, with a DGU-14A degasser, one LC-10AT pump, and a FCV-10AL Low Pressure Valve Module. Injection was through a 10 μ L injection loop. Separations were obtained on a reverse phase analytical column (Luna Omega 5 μ m Polar C18 100 LC column 150 \times 4.6 mm with a 4.0 \times 3.0 mm guard column) obtained from Phenomenex which was housed in a Shimadzu CTO-10AS column oven set to 30 $^{\circ}$ C.

A 20 minute long, two-solvent gradient method was used with 20 mM ammonium formate buffer, pH 3.20 (solvent A) and HPLC-grade acetonitrile (solvent B). The gradient steps began with 60% solvent B increased to 95% solvent B over a nine-minute period, held at 95% for three minutes and then reduced back to 60% solvent B by minute sixteen and held at 60% until the end of the run. Flow rate was maintained at

1.2 mL min⁻¹. The diode array detector was set to scan in 1 nm increments from 190 – 800 nm.

CBDA and CBD were identified by comparing their retention times to those of reference standards. Under these conditions, the retention times of CBDA and CBD were 7.0 and 8.1 minutes, respectively. Chromatograms were generated with CLASS-VP 7.2.1 SP1 software.

**Preparative High Performance Liquid
Chromatographic Method for
Purifying Cannabidiolic
Acid from a Crude
Cannabis Extract**

A Shimadzu VP HPLC was used to develop a preparative method for purifying CBDA from a crude *Cannabis* extract. The HPLC was equipped with a CBM-10AW communications bus module and a SPD-10A UV-VIS detector DGU-14A degasser, two LC-10AT pumps, and a 1.00 mL injection loop. Separations were obtained on a preparative reverse phase column (Luna Omega 5 µm Polar C18 100 LC column 150×21.2 mm with a Polar C18 SecurityGuard prep cartridge, 15×21.2 mm) obtained from Phenomenex, housed in a Shimadzu CTO-10A column oven set to 30 °C. A Gilson FC-100 microfractionator was connected to the HPLC effluent outlet and used to collect the separated compounds.

A 90 minute long, two-solvent gradient method was used with 20 mM ammonium formate buffer at 3.20 pH (solvent A) and HPLC-grade acetonitrile (solvent B). The gradient steps began with 50% solvent B held for five minutes and increased to 95% solvent B over a 55-minute period, 95% solvent B was maintained for 10 minutes, and then decreased back to 50% over the remaining 20-minute period. Flow rate was

maintained at 4.5 mL min^{-1} . The two channels of the UV-VIS detector were set to 220 nm and 274 nm. Under these conditions, the retention time of CBDA on the preparatory column was found to be 46 minutes. Chromatograms were generated with CLASS-VP 7.2.1 SP1 software.

CBDA was identified by collecting major peaks using the Gilson FC-100 microfractionator set to collect in one-minute increments beginning 40 minutes into the run. The test tubes containing the major peaks were collected from the fractionator. Test tubes which contained fractions of the same peak were combined. Each sample representing a peak in the chromatogram was placed into a 50-mL round-bottom flask. Cannabinoids were recovered from the HPLC solvent solution by first rotary evaporating the acetonitrile and then back extracting the cannabinoids from the remaining ammonium formate solution into 5.0 mL pentane. The pentane layer was transferred with a Pasteur pipette to a test tube. The pentane was evaporated to dryness using nitrogen gas and the resulting residue was resuspended in 1.0 mL ethanol.

Infrared Spectroscopy Procedure

A Thermo Scientific Nicolet iS5 IR equipped with an iD5 ATR accessory and zinc selenide crystal was used to acquire IR spectra using the OMNIC software. A 1.0 mg sample of CBDA recovered from the preparative HPLC was applied to the surface of the crystal. After the pressure arm was lowered and secured, the sample was analyzed. Obtained spectra were compared to standard spectra from a study by Hazekamp, Peltenburg, Verpoorte & Giroud (2005).

¹H- and ¹³C Nuclear Magnetic Resonance Procedure

A Bruker 400 MHz NMR was used with TOPSPIN 1.3 software program to generate all NMR spectra. Confirmatory ¹H- and ¹³C-NMR spectra were acquired for cannabidiolic acid isolated using the outlined preparative HPLC method. Since the recovered CBDA was initially dissolved in 1.0 mL ethanol, the sample was evaporated to dryness with nitrogen gas and the residue (3.7 mg) was resuspended in 600 µL of deuterated chloroform for NMR analysis. Obtained spectra were compared to standard spectra of cannabinoids from a study by Choi et al. (2004).

CHAPTER IV

RESULTS AND DISCUSSION

This study had a dual purpose, to isolate CBDA from the plant *Cannabis sativa* and to identify a microbial cell line which possesses a decarboxylase enzyme capable of converting CBDA to CBD. Before various cells could be screened for decarboxylase activity, the optimal conditions for the chromatographic column and mobile phase had to be confirmed to ensure optimal separation and identification of the cannabinoids. This analytical method was employed to monitor any enzymatic conversion of CBDA to CBD in the enzymatic assays and as a confirmatory method for preparatory isolation of CBDA. Once developed, cell lines could be monitored for CBDA decarboxylase activity. The following sections discuss results from HPLC methodology development, CBDA isolation, and enzymatic assays.

Analytical Method for Monitoring Conversion of Cannabidiolic Acid to Cannabidiol

Model compounds from previous studies were separated by HPLC on a Tosohaus C₁₈ column using an isocratic method consisting of 35% methanol and 65% 0.5 M potassium acetate buffer, pH 3.0, with a flow rate of 1.0 mL/min at room-temperature (Cribbs, 2014); however, this method was not suitable since it was not able to baseline resolve CBDA and CBD well. Separating these two compounds was particularly challenging because there is such a small difference in the chemical formula, molecular

weight, and the physical and chemical properties of CBDA and CBD. The reverse phase polar C18 (Luna Omega 5 μm Polar C18 100 Å, LC Column 150 x 4.6 mm) column was obtained from Phenomenex who suggested an isocratic mobile phase of 25% water + 0.1% formic acid and 75% acetonitrile + 0.1% formic acid reported in their applications. They also reported a fifteen minute-long gradient method using 20 mM ammonium formate, pH 3.2 (solvent A) and acetonitrile (solvent B) for a similar column with different dimensions (Luna Omega 1.6 μm C18 100 Å, LC Column 100 x 2.1 mm). This method began by increasing solvent B from 60% to 95% over 12 minutes, holding for one minute, and then decreasing solvent B back to 60% during the remaining time at a flow rate of 0.4 mL min⁻¹ at 40 °C. The gradient approach was taken; however results reported by Phenomenex® could not be replicated with the difference in columns. The initial concentration of solvent B was increased from 5% to 60%, flow rate was increased to 1.2 mL min⁻¹, and column temperature was decreased to 30 °C. This achieved greater separation of various cannabinoids. The optimized analytical method is described under the Instrumentation section in Chapter III (Methods).

With the optimized analytical HPLC method, standards of CBDA and CBD were analyzed to confirm the identity of analyte peaks from enzymatic assays and CBDA isolation by matching their retention times with that of these standards. The retention times of CBDA and CBD were determined to be 7.0 min (Figure 15) and 8.3 min (Figure 16), respectively. Baseline separation of both compounds was achieved (Figure 17) which allows for the conversion of CBDA to CBD to be monitored. Interesting to note is the difference in retention times of CBDA and CBD when in a mixture than when pure, possibly due to intermolecular interactions such as hydrogen bonding. Cannabidiolic

acid had a retention time of 7.0 min. when pure in an ethanolic solution compared to 7.5 min in an ethanolic solution with CBD. Cannabidiol's retention time in a mixture with CBDA was shifted to 8.3 min from 8.1 min when analyzed with this method.




Figure 15: Chromatogram of 2.0 mg/mL CBDA standard using the analytical HPLC method. Retention time 7.0 minutes.




Figure 16: Chromatogram of 2.0 mg/mL CBD standard using the analytical HPLC method. Retention time 8.1 minutes.




Figure 17: Chromatogram of a mixture of 1.0 mg/mL CBDA and 1.0 mg/CBD using the analytical HPLC method. CBDA retention time 7.5 min, CBD retention time 8.3 min.

Standard curves for both CBDA (Figure 18) and CBD (Figure 19) were constructed to quantify any enzymatic conversion during cell assays based qualitatively on the appearance of the CBD peak. Solutions of CBDA and CBD were prepared in triplicate over a range of concentrations (0.10 – 2.0 mg/mL) and analyzed using the analytical HPLC method. The area under the curve for each run of the same concentration was averaged and plotted against the concentration. Areas and averages for peaks from various concentrations of CBDA and CBD standards are given in Table 1 and Table 2, respectively.

Table 1: Area of Cannabidiolic Acid Peak at Different Concentrations Using the Analytical HPLC Method.

CBDA Concentration (mg/mL)	Trail 1	Trial 2	Trial 3	Average	Standard Deviation
2.0	4.07×10^7	3.90×10^7	4.00×10^7	3.99×10^7	6.96×10^5
1.5	2.92×10^7	3.46×10^7	3.11×10^7	3.16×10^7	2.27×10^6
1.0	2.48×10^7	2.26×10^7	2.08×10^7	2.27×10^7	1.64×10^6
0.5	9.26×10^6	8.30×10^6	8.22×10^6	8.59×10^6	4.72×10^5
0.1	4.72×10^6	1.06×10^6	5.36×10^6	3.71×10^6	1.89×10^6




Figure 18 : Standard calibration curve for cannabidiolic acid.

Table 2: Area of Cannabidiol Peak at Different Concentrations Using the Analytical HPLC Method.

CBD Concentration (mg/mL)	Trial 1	Trial 2	Trial 3	Average	Standard Deviation
2.0	2.40×10^7	2.48×10^7	2.38×10^7	2.42×10^7	4.56×10^5
1.5	1.87×10^7	1.87×10^7	1.93×10^7	1.89×10^7	2.42×10^5
1.0	1.70×10^7	1.67×10^7	1.52×10^7	1.63×10^7	7.79×10^5
0.5	7.94×10^6	8.07×10^6	8.02×10^6	8.01×10^6	5.29×10^4
0.1	2.05×10^6	2.15×10^6	1.94×10^6	2.05×10^6	8.57×10^4




Figure 19: Standard calibration curve for cannabidiol.

Isolation and Characterization of Cannabidiolic Acid

Preparative High Performance Liquid Chromatographic Method for Purifying Cannabidiolic Acid from a Crude Cannabis Extract

In addition to the analytical HPLC method, a preparatory HPLC method for cannabinoids was developed for the purpose of isolating pure CBDA from the crude Soxhlet *Cannabis* extract. Crude extract was prepared from 5.00 g of *Cannabis* containing a reported 6.5% CBDA (0.325 g) as outlined in Methodology. The extract was initially analyzed on the preparatory column using a two-solvent gradient, referred to as Gradient 1 (Figure 20), at 30 °C, with a flow rate maintained at 5 mL min⁻¹, however poor separation was achieved (Figure 21). The length of the gradient step from 25% to 95% acetonitrile was increased from 65 minutes to 95 minutes, while flow rate was decreased to 4.5 mL min⁻¹ to make Gradient 2 (Figure 22). This method was able to achieve separation of at least 4 major components (Figure 23).

Because the first compounds from the extract eluted one hour into the run, analysis time was shortened by 60 minutes by increasing the starting percentage of the eluent from 25% to 50% acetonitrile. The temperature and flow rate from Gradient 2 was maintained. The resulting method (Gradient 3, Figure 24) was able to resolve cannabinoids (Figure 25) in a similar manner to Gradient 2 in 30 minutes less and with 32.5% less solvent than the originally suggested method using Gradient 1. Both time and materials can be costly so optimizing the method in this way was ideal to save valuable

resources with very simple adjustments. The optimized preparatory method is described under “Instrumentation” in Chapter III (Methods).




Figure 20: Gradient Method 1




Figure 21: Chromatogram of CBDA extract from Soxhlet extraction using the preparatory HPLC method with Gradient 1.




Figure 22: Gradient Method 2




Figure 23: Chromatogram of CBDA extract from Soxhlet extraction using the preparatory HPLC method with Gradient 2.




Figure 24: Gradient Method 3




Figure 25: Chromatogram of CBDA extract the from Soxhlet extraction using the preparatory HPLC method with Gradient 3.

When the sample concentration exceeds the range of the detector, the detector cannot generate a larger response, and typically a flat-topped peak is observed, referred to as detector overload (Dolan, 2015). Two flat-topped peaks in the chromatogram were observed, so to address this issue the extract was diluted with two successive 1:2 dilutions with absolute ethanol prior to injection on the HPLC which was sufficiently dilute to prevent detector overload (Figure 26). This dilution factor also allowed for only the seven most abundant cannabinoids to be detected.




Figure 26: Chromatogram of CBDA extract, after two successive 1:2 dilutions with absolute ethanol from the Soxhlet extraction using Gradient 3.

Chromatographic Characterization of Cannabidiolic Acid

Once the preparative HPLC method was optimized, fractions were collected every minute to isolate each individual peak for analysis. Thirty test tubes of 4.5 mL of eluent were collected in total. Of the thirty, nine test tubes contained compounds of interest which corresponded to seven peaks on the chromatogram and these were collected for

analysis. A summary of fractions collected is given in Table 3. Fractions were prepared for analysis by combining test tubes which contained eluent from the same peak. The acetonitrile was removed by rotary evaporation and the cannabinoids were extracted from the remaining ammonium formate solution with 5 mL of pentane. The pentane was then evaporated to dryness with nitrogen gas; the resulting residue was resuspended in 1.0 mL absolute ethanol and analyzed by the analytical HPLC method.

Table 3: Summary of Collected Fractions from the Preparatory HPLC Method.

Collection Time	Fraction Number	Peak Number (Figure 26)	Peak Area	Retention Time (min.)	Identity
40-44	1-5	N/A	-	-	-
45-46	6-7	1	251815642	46.025	CBDA
47	8	2	53227248	47.683	Unknown
48-49	9-10	N/A	-	-	-
50	11	3	75508734	50.492	Unknown
51	12	N/A	-	-	-
52	13	4	16942665	52.300	Unknown
53-55	14-16	N/A	-	-	-
56	17	5	17919678	56.700	Unknown
57-58	18, 19	N/A	-	-	-
59	20	6	17791656	59.233	Unknown
60-61	21, 22	7	294579036	60.758	THCA
62-70	23-30	N/A	-	-	-

The most abundant cannabinoids of *Cannabis sativa* are Δ^9 -THCA and CBDA (Elsohly & Slade, 2005). The area under the curve of peak 1 and peak 7 accounted for more than 55% of all compounds detected, making up 29.92% and 25.58 % of the area, respectively as reported in the generated area report. Because peak 1 and 7 (Figure 26) were the most abundant, this suggested that the identity of the compounds which produced these peaks may be CBDA, the analyte of interest, and Δ^9 -THCA, the other most abundant cannabinoid in the extracted *Cannabis* sample. The compound from the

first major peak, collected from fractions 6 and 7, had a retention time of 46.025 minutes using the preparatory HPLC method. When analyzed with the analytical HPLC method, the compound had a retention time of 7.0 minutes (Figure 27) which matched the retention time for the standard CBDA (Figure 28). This was the first indication that CBDA had been successfully recovered from the Soxhlet extraction of the *Cannabis* plant material followed by the preparatory HPLC method. The identity of the compound which produced peak 1 (Figure 26) as CBDA was further supported by NMR and IR spectroscopy.




Figure 27: Chromatogram using the analytical HPLC method of the compound recovered from peak 1. Retention time 7.0 minutes.




Figure 28: CBDA standard (2.0 mg/mL) using the analytical HPLC method. Retention time 7.0 minutes.

Nuclear Magnetic Resonance Spectral Analysis

Both ^1H -NMR and ^{13}C -NMR spectra of the compound from peak 1 were obtained for comparison to spectra reported in the literature by Choi et al. (2004) and by Hazekamp, Choi and Verpoorte (2004). The study by Choi et al. (2004), which studied Δ^9 -THCA, THC, Δ^8 -THC, CBG, CBN, CBDA, CBD, and two flavonoids, was used for overall assignment of each peak in the ^1H -NMR and ^{13}C -NMR spectra. Unique singlets in the range of δ 4.0—7.0 in the ^1H -NMR spectrum of Δ^9 -THCA, THC, CBDA, CBD, and CBN reported by Hazekamp et al., were used to further support the identity of CBDA once the spectrum was fully assigned. The numbering scheme for cannabidiolic acid used in these two studies is given in Figure 29. Although the spectra for cannabidiolic acid from the Choi et al. study were not provided, they reported a table of chemical shifts (ppm) with reference to TMS. Table 7 compares actual values from the ^1H - and ^{13}C -NMR obtained for the compound in peak 1 with values reported in the Choi et al. study for cannabidiolic acid.




Figure 29: Numbering scheme used by Choi et al. (2004) for CBDA.

Table 4: Comparison of ^1H - and ^{13}C -NMR Spectra of the Compound Isolated from Peak 1 to Assignments for Cannabidiolic Acid Reported by Choi et al. (2004).

Position	^1H NMR		^{13}C NMR	
	ppm (actual, peak 1)	ppm (literature)	ppm (actual)	ppm (literature)
1	4.09	4.09	35.36	37.3
2	5.57	5.56	123.84	126.9
3			140.55	133
4	2.23	2.2	30.17	31.7
5	*	1.86	27.7	30.8
6	2.38	2.4	46.6	45.7
7	1.8	1.8	23.7	23.6
8			147.8	150.4
9	4.54	4.55	111.34	110.5
	4.39	4.4		
10	1.72	1.72		
1'			18.86	19.3
2'			114.45	116.5
3'			164.1	161.8
4'			102.5	104.2
5'	6.26	6.26	147.1	147
6'			111.9	111.3
1''	2.92, 2.82	2.92, 2.82	161	150.4
2''	1.58	1.57	36.56	37.6
3''	1.33	1.33	31.2	32.8
4''	1.33	1.33	31.97	33.2
5''	0.89	0.89	22.54	23.5
2'-OH	11.85	11.93	14.07	14.4
6'-OH	6.65	6.63		
COOH			176.06	175.6

*peak is absent in the spectra, assumed to be buried beneath the signal at 1.8 ppm

In the ^{13}C -NMR spectrum of the isolated compound (Figure 30), twenty-two distinct carbon peaks were observed as would be expected for cannabidiolic acid ($\text{C}_{22}\text{H}_{30}\text{O}_4$) (Figure 29). Choi et al. reported only twenty-one distinct peaks, noting that carbons in position 8 and 6' overlapped and appear as one signal at 150.4 ppm. Differences in chemical shifts from those reported in the literature could be attributed to the fact that Choi et al. obtained ^{13}C NMR spectra at 100 MHz in deuterated methanol, whereas the spectra obtained in this study were obtained at 100 MHz in deuterated

chloroform making it possible for the electronic environment of the atoms in these solutions to be different due to solute-solvent interactions.




Figure 30: ^{13}C -NMR spectrum of the compound isolated from peak 1 in CDCl_3 .

Unlike the ^{13}C -NMR, conditions for obtaining ^1H -NMR spectra were the same in both the literature references and in this study. As shown in Figure 31, a carboxylic acid proton is observed at 11.858 ppm, as would be expected from a carboxylic acid proton. Choi et al. reported the proton signal at 11.93 is from the 6' -OH but did not report a carboxylic acid hydrogen which typically is observed between 10-13 ppm. Phenolic hydrogens with intramolecular hydrogen bonding have a much broader range (10.5-16 ppm) (Jackman and Sternhell, 1969), thus it is possible that the signal is indeed from the 6'-OH and that the carboxylic acid hydrogen is absent. A carboxylic acid carbon signal can be observed, however, in the ^{13}C spectrum at 176 ppm, thus it can be confirmed that

the carboxylic acid carbon is present. The H-5 signal was not observed in the spectrum for the compound from peak 1, however the H-7 peak at 1.80 ppm integrated for six hydrogens when it should integrate for three hydrogens. The H-5 signal that is supposed to be present at 1.86 ppm could be buried beneath this peak, accounting for two hydrogens, suggesting that the last hydrogen which was unaccounted for in the integration to be the 6'-OH hydrogen.




Figure 31: ^1H -NMR spectrum of compound isolated from peak 1 in CDCl_3

Because the disputed signals are from acidic hydrogens in CBDA, a technique for confirming signals from exchangeable protons called a deuterium exchange was performed by adding 100 μL of D_2O directly into the NMR tube containing the sample. Spectra of the CBDA isolated from the preparatory column before (A) and after (B) the addition of D_2O are presented in Figure 32. The absence of the signals at 11.85 and 6.65 ppm in the spectrum after deuterium exchange (Fig. 32 B) confirms that those two signals came from exchangeable hydrogens. No additional information was gained as to the assignments of the carboxylic acid -OH, 6'-OH, and 2'-OH. In order to further validate that the compound was in fact CBDA, signals in the 4.0-7.0 region where distinguishable

singlets between cannabinoids can be observed were compared. Four distinct signals for H-4, H-10, H-9 *trans* and H-10 *cis* are reported at 6.26, 5.55, 4.54, and 4.40 in the literature (Hazekamp et al., 2004) and observed in the spectrum obtained.




Figure 32: Deuterium exchange experiment for the compound isolated from peak 1. (A) before the addition of D₂O and (B) after the addition of D₂O.

Infrared Spectral Analysis

Additional spectral analysis included infrared spectroscopy which was obtained from a viscous 1.0 mg sample of the compound isolated from peak 1. A broad peak in the IR spectrum, starting at approximately 3400 cm⁻¹, can be observed (Figure 33) as well as a sharp peak at 1617.30 cm⁻¹ and a sharp peak at 1257.32 cm⁻¹. These three signals together are indicative of a carboxylic acid -OH and its C=O and C-O stretches.

Comparing this spectrum to the IR spectrum by Hazekamp et al. (2005), both have a broad carboxylic acid peak, along with sharp peaks at approximately 1250 cm⁻¹. The peaks protruding around 2900 cm⁻¹ match C-H stretched indicated by Hazekamp et al.

This spectrum provided further confirmation that the identity of the compound present in peak 1 is CBDA.




Figure 33: IR analysis of the compound isolated from peak 1.

Recovery

The *Cannabis* plant material contained a reported 6.5% CBDA. To determine the percent of CBDA recovered by preparatory HPLC following Soxhlet extraction, a 0.5 mL sample of pure extract was separated by preparatory HPLC, collected by the fractionator, and processed with rotary evaporation and back-extraction into pentane. The pentane was evaporated in a pre-weighed container and the mass of the recovered CBDA was determined for the 0.5 mL injection volume. This mass was multiplied by 32 to account for the total volume of 16 mL for the extract. On average, 77.4 % of the reported CBDA was recovered from the 0.325 g expected from 5.00 g plant material. Table 5 summarizes recovery data.

Table 5: Percent Recovery Data

Injection	CBDA Recovered* (g)	Calculated total CBDA extracted (g)	% CBDA Recovered
1	0.00755	0.2416	74.34
2	0.00841	0.2691	82.81
3	0.00697	0.2440	75.08

*Recovered from 0.5 mL injection of pure extract

Other Cannabinoid Analyses

Although CBDA was the focus of this study, this preparatory method could potentially be used to separate a variety of compounds extracted from *Cannabis* plant material. The compound from the last major peak, peak 7, collected from fractions 21 and 22, had a retention time of 60.8 minutes on the preparatory column. When analyzed with the analytical HPLC method, the compound had a retention time of 10.8 minutes (Figure 34) which was comparable with a reference standard for THCA with a retention time of 10.6 min (Figure 35). When THCA is in a mixture of cannabinoids, it appears to elute slightly later than when in an pure solution, similar to the effect observed when CBDA and CBD are in a mixture.




Figure 34: Chromatogram using the analytical HPLC method of the compound recovered from peak 7 using the preparatory HPLC method. Retention time 10.8 minutes.




Figure 35: Chromatogram of THCA standard (1.0 mg/mL). Retention time 10.6 minutes

In an effort to identify which other cannabinoids were present, an additional preparatory run was performed but instead of collecting individual fractions all compounds which eluted between 40 and 70 minutes were collected. This fraction was prepared for analysis and chromatographed on the analytical column (Figure 36), which was compared to the phytocannabinoid standard mixture (Figure 37) that contained 100 $\mu\text{g/mL}$ of each of the following cannabinoids: CBDA, cannabigerolic acid (CBGA), cannabigerol (CBG), CBD, tetrahydrocannabivarin (THCV), cannabinol (CBN), Δ^9 -tetrahydrocannabinolic acid (Δ^9 -THCA), Δ^9 -tetrahydrocannabinol (Δ^9 -THC), Δ^8 -tetrahydrocannabinol (Δ^8 -THC), and cannabichromene (CBC).




Figure 36: Chromatogram generated using the analytical HPLC method of the cannabinoid mixture recovered from the 40-70 minute fraction using the preparatory HPLC method with Gradient 3.




Figure 37: Chromatogram of phytocannabinoid mixture standard using the analytical HPLC method.

It appears that, like CBDA and CBD, other cannabinoids will elute at different times in a mixture possibly due to interactions between compounds than when in a pure sample. It was assumed that the peak at 7.9 minutes (Figure 36) is CBDA and the peak at 11.3 minutes is THCA due to abundance and order of elution. It appears that the crude extract contains several of the standard cannabinoids: one which elutes at 8.4 min., the next at 8.9 min., followed by one at 9.6 min. and one which elutes at 10.1 min. Each fraction was analyzed with the analytical HPLC method and chromatograms were collected (Appendix B). These chromatograms indicated that each sample was impure and therefore not suitable for further analysis to confirm the identity. If pure cannabinoid standards were obtained, other than the major cannabinoids, more studies could follow to identify which of the standards match the compounds in this crude extract; however, this was outside the scope of this study.

Cell Assay for Decarboxylase Activity

To identify a microbial species which exhibits decarboxylase activity, seven cell lines were screened, four bacterial and three fungal. Species were chosen based upon

literature reports of decarboxylase activity for compounds similar in structure to CBDA. Species were screened using cells which were either intact or lysed and either induced or uninduced depending on the assay. A summary of cell assays is given in Table 6. A full description of the various combinations of conditions for each assay is given in Table 7 (Appendix A).

Table 6: Summary of Cell Incubations for Decarboxylase Activity.

Cell Type	Intact	Lysed	Induced	Induction Substrate
<i>Aspergillus niger</i>	No	Yes	Yes	Salicylic acid
<i>Aspergillus clavatus</i>	No	Yes	Yes	Salicylic acid
<i>Bacillus subtilis</i>	Yes	Yes	Yes	Salicylic acid or 4-hydroxybenzoic acid
<i>Enterobacter (Klebsiella) aerogenes</i>	Yes	Yes	No	Not induced
<i>Klebsiella pneumonia</i>	Yes	Yes	Yes	2,4-dihydroxybenzoic acid
<i>Penicillium chrysogenum</i>	No	Yes	Yes	Salicylic acid
<i>Pseudomonas putida</i>	Yes	Yes	Yes	Salicylic acid or 2,4-dihydroxybenzoic acid

From the bacteria screened, *B. subtilis*, *E. aerogenes*, and *K. pneumonia* displayed no detectable CBDA decarboxylase activity; representative chromatograms for each species are given in Appendix C1, C2, and C3, respectively. *B. subtilis* was assayed under various combinations of intact/lysed and uninduced/induced conditions, however no CBDA decarboxylase activity was observed. This cell line was not tested with model compounds but was chosen because it expresses the genes for both a phenolic acid decarboxylase and a phenylacrylic acid decarboxylase. Duy et al. (2007) showed that these enzymes were both expressed when induced with salicylic acid. However, even

after induction with salicylic acid no CBDA decarboxylase activity was observed in the current study when *B. subtilis* was incubated in the presence of CBDA.

E. aerogenes contains an enzyme which decarboxylates both Category 1 and 2 substrates, and this cell line was assayed both intact and lysed. Earlier work by Cribbs (2014) demonstrated the ability of lysed *E. aerogenes* cells to decarboxylate the Category 2 substrate p-hydroxybenzoic acid but not any phenolic carboxylic acid with hydroxyl groups *ortho* to the carboxylic acid. A permeability barrier for the substrate of the decarboxylase was noted in the study by Grant and Patel (1969) which could explain the lack of activity in intact cells. An optimal pH of 6 was also noted which could explain a lack of detectable activity when incubated in the presence of CBDA, even with lysed cells, because all assays were performed at the pH of the media.

Both intact and lysed *K. pneumonia* cells, induced with 2,4-dihydroxybenzoic acid, were incubated in the presence of CBDA. No decarboxylase activity was observed in any of the three assays performed. Like *E. aerogenes*, the decarboxylase enzyme from *K. pneumonia* is able to decarboxylate the Category 2 substrate p-hydroxybenzoic but no model compound with a phenolic hydroxyl group *ortho* to the carboxylic acid (Cribbs, 2014).

None of the fungal species screened displayed CBDA decarboxylase activity when incubated in the presence of CBDA. Representative chromatograms for assays of *A. clavatus*, *A. niger*, and *P. chrysogenum* are given in Appendix C4, C5, and C6, respectively. *Aspergillus clavatus* was not tested with model compounds but was chosen because it has the *patG* gene, which codes for an enzyme that decarboxylates 6-methylsalicylic acid. Two assays of *Aspergillus clavatus* induced with salicylic acid and

one assay of *A. clavatus* induced with CBDA were performed but no CBDA decarboxylase activity was observed. *Penicillium chrysogenum* and *Aspergillus niger* did not demonstrate any decarboxylase activity when incubated with model compounds (Cribbs, 2014) or with CBDA in the current study.

In previous studies, *Pseudomonas putida* was demonstrated to catalyze the decarboxylation of 2-hydroxy-4-methoxybenzoic acid (Cribbs, 2014). In the current study, twelve assays were performed for CBDA decarboxylase activity with *P. putida*. A representative set of chromatograms are presented in Figures 38-42. Individual treatments for each assay are given in Table 7, (Appendix A) and additional representative chromatograms are given in Appendix C7. CBDA typically elutes at 7.0 minutes with the analytical HPLC method; however the column oven was not set, and samples were run at ambient room temperature for this assay, resulting in retention times of the standards for CBDA and CBD 7.8 and 8.3 minutes, respectively. Figure 38 shows a chromatogram of an incubation extraction at time 0 with only one major peak, CBDA at 7.8 minutes. After 1 hour, CBDA was still the only species present (Figure 39). After 2 hours of incubation, the peak for CBDA was still be observed, and a peak for CBD appeared at 8.3 minutes (Figure 40). The observation of both peaks persists at time 3 hr. (Figure 41) and 4 hr. (Figure 42), with a slight increase in peak intensity in the final hour, t = 4 hr., of assay.




Figure 38: Chromatogram of CBDA after 0 hours of incubation with lysed *P. putida*. CBDA is observed at 7.8 min.




Figure 39: Chromatogram of CBDA after 1 hour of incubation with lysed *P. putida*. CBDA is observed at 7.8 min.




Figure 40: Chromatogram of CBDA after 2 hours of incubation with lysed *P. putida*. CBDA (7.8 min) and CBD (8.3 min) are observed




Figure 41: Chromatogram of CBDA after 3 hours of incubation with lysed *P. putida*. CBDA (7.8 min) and CBD (8.3 min) are observed.




Figure 42: Chromatogram of CBDA after 4 hours of incubation with lysed *P. putida*. CBDA (7.8) and CBD (8.3) are observed.

An autoclaved sample of the *P. putida* used as a control was assessed concurrently with the active cell culture to be sure any observed conversion was, in fact, produced by an enzyme present in the cell. Only CBDA was observed for all time points with the control. The observation of the appearance of CBD suggests that *Pseudomonas putida* has an enzyme which can convert the prodrug CBDA into the physiologically active CBD. To calculate the rate at which this occurs, the area under the curve for CBD was taken and calculations were performed in Excel using the volume of the injection (10 μL) and the molar mass of CBD (314.46 g/mol). The rate at which CBD appeared was determined to be 5.0 nmol/hour.

CHAPTER V

CONCLUSIONS

The first goal of this research was to isolate and identify CBDA from an extract prepared in-house from *Cannabis* plant material. Isolating cannabinoids in-house from plant material has several benefits for this research. The plant material itself is considerably less expensive relative to the cost of a pure, isolated cannabinoid. Reagent cost and a minimal on-hand quantity of the desired cannabinoid can hinder productive research, thus having the ability to isolate CBDA lowers cost and increases efficiency. In-house isolation also allows a more “green” approach because one is not required to use extra resources to ship reagents and can monitor quality control.

A Soxhlet extraction was performed to obtain an extract which was high in CBDA. The extract was then purified using a preparative HPLC method developed for the separation of cannabinoids. The CBDA obtained from this method was seemingly pure, as no other peaks were observed in the analytical HPLC chromatogram obtained and few unidentifiable peaks were observed in the ^1H - and ^{13}C - NMR spectra. Other cannabinoids could possibly be isolated with this method as well, however, CBDA was the only pure cannabinoid isolated in this study. The development of this method will allow further research to be conducted in our laboratory on various independently isolated cannabinoids.

The second goal of this research was to identify a cell line possessing enzymes capable of decarboxylating CBDA. In its physiologically active form, cannabidiol is supported by a vast body of literature as a potential cancer treatment. To accomplish this goal, several cell lines were screened for CBDA decarboxylase activity by monitoring conversion of CBDA to CBD via HPLC. An analytical HPLC method was successfully developed for this task. Cell lines were selected based on the literature and previous work in our research group with compounds which model cannabinoids (Cribbs, 2014).

The results indicate that a CBDA decarboxylase is present in *Pseudomonas putida*. Cultures of *P. putida* which were not induced or induced with 2,4-dihydroxybenzoate did not show any CBDA decarboxylase activity. However, cultures of *P. putida* that were induced with salicylic acid for three days and lysed prior to decarboxylase activity screening indicate the formation of CBD when incubated with CBDA for at least three hours. This decarboxylase activity from *P. putida* is the first enzymatic method for the decarboxylation of CBDA observed which does not require substantial heat to convert it to its physiologically active form CBD.

If a cell line was identified, therapeutic drug development can move forward using Cell-in-a-Box® technology. With the desired decarboxylation of CBDA to CBD observed, isolation of the gene for the decarboxylation enzyme would be the next logical step with cultures of *P. putida* which have been induced for three days with salicylic acid and lysed. The isolated gene could then be inserted into a human embryonic kidney (HEK) cell through genetic engineering. Once the HEK cell has been transfected with the decarboxylase gene, it can be analyzed with the same assay developed for microbial cells tested in this study, with the ultimate goal of encapsulating the HEK cells using

Cell-in-a-Box® technology and testing in clinical trials. This could be especially beneficial for patients with tumors which are difficult to remove surgically such as gliomas, which CBD has been shown to treat effectively. The specific activity should also be quantified because literature studies show that the effectiveness of CBD is dose-dependent.

Future Research

The preparatory HPLC method could be developed further to not only isolate CBDA but other major cannabinoids as well. The lack of purity of the other major cannabinoids collected in this study meant that obtaining NMR data would not prove useful for identification in this study. If the samples were pure, these could be collected and used to identify compounds other than CBDA and THCA. Current standards could also be spiked with standards to help identify unknown peaks in the current chromatograms.

The procedure for collecting fractions would have to be optimized in order to obtain compounds which are purer than those obtained in this study. This may be achieved by reducing the time increment for collection or by altering the preparatory HPLC parameters to achieve greater baseline separation between peaks. This could be achieved by using a different organic component to manipulate the polarity of the mobile phase. Acetonitrile could be substituted by a more polar solvent such as methanol which may increase run length but could provide better resolution of peaks. Methanol is also generally less expensive than acetonitrile with current prices from Sigma-Aldrich quoted at \$69.50 and \$223.00, respectively, for 1 L of HPLC-grade reagents.

The percent recovery of cannabidiolic acid could also be improved by minimizing transfer steps in the sample processing once collected from the HPLC. The current process requires evaporating the organic solvent and then back extracting from the aqueous phase into pentane. If the number of steps in this process were able to be reduced perhaps less CBDA would be lost in transfer and could increase the percent recovery.

Additional decarboxylase assays could be done to find additional microbial species capable of the decarboxylation of CBDA to CBD, with a focus on fungal species to determine if there might be a species which catalyzes this reaction more efficiently. The decarboxylase expressed by *A. clavatus* decarboxylates Category 1 substrates, those which are the most similar to the structure of CBDA; however, there was difficulty in homogenizing fungal samples when preparing them for assay. Work should be done with current collaborators who prepare cell cultures to optimize this method and then additional assays should be performed with cultures of different *Aspergillus* species such as *A. clavatus*, *A. giganteus*, and *A. longivesica* which were reported by Snini et al. (2013) to express the *PatG* gene. Another fungus which was not tested was *Trichosporon moniliform* which decarboxylates salicylic acid and many other carboxylic acids (Kirimura et al., 2010). Care needs to be taken in the initial stages of working with these types of cells since there is a risk of fungal infection. The risks associated with these types of cells should be mitigated by incorporating the gene for the decarboxylase in a genetically-engineered human embryonic kidney cell.

If additional decarboxylase assays are in order, a standard curve should be made in which CBDA and CBD are analyzed in tandem. This would assure that any effects

observed from interaction of each analyte with each other could be observed. This would also be a better application since they are not being analyzed individually in the context of the enzymatic assay. A standard curve which matches the application would be more beneficial than the current format of each individual standard curve.

Advancement in understanding of the therapeutic benefits of cannabinoids has propelled the scientific community forward to develop a multitude of cannabinoid-based therapies. A significant hurdle on the road to making the most beneficial medicine available to the patients has been delivering the appropriate dose of cannabinoids in a safe and effective manner. The results from this research provide an alternative to the traditional mode of activating cannabinoid pro-drugs and a new way to potentially control and monitor the effective dose of cannabinoids being administered.

References

- Alexander, A., Smith, P. F., & Rosengren, R. J. (2009). Cannabinoids in the treatment of cancer. *Cancer Letters*, 285(1), 6-12.
- Alger, B. E. (2013). Getting High on the Endocannabinoid System. *Cerebrum*, 2013(14).
- Alvares, L. D., Genro, B. P., Diehl, F., & Quillfeldt, J. A. (2008). Differential role of the hippocampal endocannabinoid system in the memory consolidation and retrieval mechanisms. *Neurobiology of Learning and Memory*, 90(1), 1-9.
- Backer, B. D., Debrus, B., Lebrun, P., Theunis, L., Dubois, N., Decock, L., . . . Charlier, C. (2009). Innovative development and validation of an HPLC/DAD method for the qualitative and quantitative determination of major cannabinoids in *Cannabis* plant material. *Journal of Chromatography B*, 877(32), 4115-4124.
- Benezra, R., Davis, R., Lockshon, D., Turner, D., & Weintraub, H. (1990). The protein Id: A negative regulator of helix-loop-helix DNA binding proteins. *Trends in Genetics*, 6, 174.
- Calvaruso, G., Pellerito, O., Notaro, A., & Giuliano, M. (2012). Cannabinoid-associated cell death mechanisms in tumor models. *International Journal of Oncology*, 41(2), 407-413.
- Cancer Statistics. (2017). Retrieved from <https://www.cancer.gov/about-cancer/understanding/statistics>

- Casarett, L., & Doull, J. (2013) *Toxicology The Basic Science of Poisons*. (C. Klaassen, Ed.) (8th ed.). New York: McGraw-Hill Education.
- Cavin, J., Dartois, V., & Divies, C. (1998). Gene cloning, transcriptional analysis, purification, and characterization of phenolic acid decarboxylase from *Bacillus subtilis*. *Applied and Environmental Microbiology*, 64(4), 1466-1471.
- Cianchi, F., Papucci, L., Schiavone, N., Lulli, M., Magnelli, L., Vinci, M. C., . . . Masini, E. (2008). Cannabinoid Receptor Activation Induces Apoptosis through Tumor Necrosis Factor α -Mediated Ceramide De novo Synthesis in Colon Cancer Cells. *Clinical Cancer Research*, 14(23), 7691-7700.
- Choi, Y. H., Hazekamp, A., Peltenburg-Looman, A. M., Frédérich, M., Erkelens, C., Lefeber, A. W., & Verpoorte, R. (2004). NMR assignments of the major cannabinoids and cannabiflavonoids isolated from flowers of *Cannabis sativa*. *Phytochemical Analysis*, 15(6), 345-354.
- Cribbs, J. (2014). *Enzymatic conversion of model compounds that represent pharmacologically active pro-drugs*. (Unpublished master's thesis). University of Northern Colorado.
- Curiel, J. A., Rodríguez, H., Landete, J. M., Rivas, B. D., & Muñoz, R. (2010). Ability of *Lactobacillus brevis* strains to degrade food phenolic acids. *Food Chemistry*, 120(1), 225-229.
- Dolan, J. W. (2015). Overload in liquid chromatography. Retrieved from <http://www.chromatographyonline.com/overload-liquid-chromatography-0>

- Duy, N. V., Mäder, U., Tran, N. P., Cavin, J., Tam, L. T., Albrecht, D., . . . Antelmann, H. (2007). The proteome and transcriptome analysis of *Bacillus subtilis* in response to salicylic acid. *Proteomics*, 7(5), 698-7105
- Eisenreich, W., Rohdich, F., & Bacher, A. (2001). Deoxyxylulose phosphate pathway to terpenoids. *Trends in Plant Science*, 6(2), 78-84.
- Ellert-Miklaszewska, A., Grajkowska, W., Gabrusiewicz, K., Kaminska, B., & Konarska, L. (2007). Distinctive pattern of cannabinoid receptor type II (CB2) expression in adult and pediatric brain tumors. *Brain Research*, 1137, 161-169.
- Elphick, M. R. (2012). The evolution and comparative neurobiology of endocannabinoid signaling. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1607), 3201-3215.
- ElSohly, M., & Gul, W. (2016). *Handbook of Cannabis*. Oxford: Oxford Univ Press.
- Elsohly, M. A., & Slade, D. (2005). Chemical constituents of marijuana: The complex mixture of natural cannabinoids. *Life Sciences*, 78(5), 539-548.
- Fellermeier, M., Eisenreich, W., Bacher, A., & Zenk, M. H. (2001). Biosynthesis of cannabinoids Incorporation experiments with ¹³C-labeled glucoses. *European Journal of Biochemistry*, 268(6), 1596-1604.
- Fernández-Ruiz, J., Romero, J., Velasco, G., Tolón, R. M., Ramos, J. A., & Guzmán, M. (2007). Cannabinoid CB2 receptor: A new target for controlling neural cell survival? *Trends in Pharmacological Sciences*, 28(1), 39-45.
- Fonseca, B., Costa, M., Almada, M., Correia-Da-Silva, G., & Teixeira, N. (2013). Endogenous cannabinoids revisited: A biochemistry perspective. *Prostaglandins & Other Lipid Mediators*, 102-103, 13-30.

- Gaoni, Y., & Mechoulam, R. (1964). Isolation, structure, and partial synthesis of an active constituent of hashish. *Journal of the American Chemical Society*, 86(8), 1646-1647.
- Grant, D., & Patel, J. (1969). The non-oxidative decarboxylation of *p*-hydroxybenzoic acid, gentisic acid, protocatechuic acid and gallic acid by *Kelbsiella aerogenes* (*Enterobacter aerogenes*). *Antonie van Leeuwenhoek*, 35, 325-343.
- Guzmán, M. (2003). Cannabinoids: Potential anticancer agents. *Nature Reviews Cancer*, 3(10), 745-755.
- Guzmán, M., Duarte, M. J., Blázquez, C., Ravina, J., Rosa, M. C., Galve-Roperh, I., . . . González-Feria, L. (2006). A pilot clinical study of Δ^9 -tetrahydrocannabinol in patients with recurrent glioblastoma multiforme. *British Journal of Cancer*, 95(2), 197-203.
- GW Pharmaceuticals. (2018, June 26). About Us. Retrieved from <http://www.gwpharm.com/about-us>
- Hallak, J., Dursun, S., Bosi, D., De Macedo, L., Machado-de-Sousa, J., Abrao, J., . . . Baker, G. (2011). The interplay of cannabinoid and NMDA glutamate receptor systems in humans: Preliminary evidence of interactive effects of cannabidiol and ketamine in healthy human subjects. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 35(1), 198-202.
- Hazekamp, A., Choi, Y. H., & Verpoorte, R. (2004). Quantitative Analysis of Cannabinoids from Cannabis sativa Using ^1H -NMR. *Chemical & Pharmaceutical Bulletin*, 52(6), 718-721.

- Hazekamp, A., Peltenburg, A., Verpoorte, R., & Giroud, C. (2005). Chromatographic and Spectroscopic Data of Cannabinoids from *Cannabis sativa* L. *Journal of Liquid Chromatography & Related Technologies*, 28, 2361-2382.
- Halvorson, H. (1963). Formation of resorcinol as an intermediate in the metabolism of 2,4-dihydroxybenzoic acid and its corresponding aldehyde. *Biochemical and Biophysical Research Communications*, 10(6), 440-443.
- Hampson, R. E., & Deadwyler, S. A. (1999). Cannabinoids, hippocampal function and memory. *Life Sciences*, 65(6-7), 715-723.
- Herkenham, M., Lynn, A. B., Little, M. D., Johnson, M. R., Melvin, L. S., De Costa, B. R., & Rice, K. C. (1990). Cannabinoid receptor localization in brain. *Proc. Natl. Acad. Sci. USA*, 87, 1932-1936.
- Hua, T., Vemuri, K., Pu, M., Qu, L., Han, G., Wu, Y., . . . Liu, Z. (2016). Crystal Structure of the Human Cannabinoid Receptor CB1. *Cell Press*, 167(3), 750-762.
- Iding, H., Dünwald, T., Greiner, L., Liese, A., Müller, M., Siegert, P., . . . Pohl, M. (2000). Benzoylformate Decarboxylase from *Pseudomonas putida* as Stable Catalyst for the Synthesis of Chiral 2-Hydroxy Ketones. *Chemistry - A European Journal*, 6(8), 1483-1495.
- Izzo, A. A., Borrelli, F., Capasso, R., Di Marzo, V., & Mechoulam, R. (2009). Non-psychoactive plant cannabinoids: New therapeutic opportunities from an ancient herb. *Trends in Pharmacological Sciences*, 30(12), 609.
- Jackman, L. M., & Sternhell, S. (n.d.). *Applications of Nuclear Magnetic Resonance Spectroscopy* (2nd ed.). London: Pergamon Press.

- Jiang, W. (2005). Cannabinoids promote embryonic and adult hippocampus neurogenesis and produce anxiolytic- and antidepressant-like effects. *Journal of Clinical Investigation*, 115(11), 3104-3116.
- Jones, P. G., Falvello, L., Kennard, O., Sheldrick, G. M., & Mechoulam, R. (1977). Cannabidiol. *Acta Crystallographica Section B Structural Crystallography and Crystal Chemistry*, 33(10), 3211-3214.
- Katchan, V., David, P., & Shoenfeld, Y. (2016). Cannabinoids and autoimmune diseases: A systematic review. *Autoimmunity Reviews*, 15(6), 513-528.
- Kirimura, K., Gunji, H., Wakayama, R., Hattori, T., & Ishii, Y. (2010). Enzymatic Kolbe-Schmitt reaction to form salicylic acid from phenol: Enzymatic characterization and gene identification of a novel enzyme, *Trichosporon moniliiforme* salicylic acid decarboxylase. *Biochemical and Biophysical Research Communications*. 392(2). 279-284.
- Klein, R. F. (2015). Analysis of marijuana by liquid chromatographic techniques: A literature survey, 1990-2015. *Microgram Journal*, 12(1-4). 1-17.
- Li, H. (1974). An Archaeological and Historical Account of *Cannabis* in China. *Economic Botany*, 28(4), 437-488.
- Ligresti, A. (2006). Antitumor activity of plant cannabinoids with emphasis on the effect of cannabidiol on human breast carcinoma. *Journal of Pharmacology and Experimental Therapeutics*, 318(3), 1375-1387.
- Mahdi, J. G., Mahdi, A. J., Mahdi, A. J., & Bowen, I. D. (2006). The historical analysis of aspirin discovery, its relation to the willow tree and antiproliferative and anticancer potential. *Cell Proliferation*, 39(2), 147-155

- Marzo, V. D. (1998). 'Endocannabinoids' and other fatty acid derivatives with cannabimimetic properties: Biochemistry and possible physiopathological relevance. *Biochimica Et Biophysica Acta (BBA) - Lipids and Lipid Metabolism*, 1392(2-3), 153-175.
- Massi, P., Vaccani, A., Ceruti, S., Colombo, A., Abbracchio, M. P., & Parolaro, D. (2004). Antitumor Effects of Cannabidiol, a Nonpsychoactive Cannabinoid, on Human Glioma Cell Lines. *Journal of Pharmacology and Experimental Therapeutics*, 308(3), 838-845.
- Massi, P., Solinas, M., Cinquina, V., & Parolaro, D. (2013). Cannabidiol as potential anticancer drug. *British Journal of Clinical Pharmacology*, 75(2), 303-312.
- Matsuda, L. A., Lolait, S. J., Brownstein, M. J., Young, A. C., & Bonner, T. I. (1990). Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature*, 346(6284), 561-564.
- Mavromoustakos, T., Yang, D., & Makriyannis, A. (1995). Small angle X-ray diffraction and differential scanning calorimetric studies on O-methyl-(Δ^8 -tetrahydrocannabinol and its 5' iodinated derivative in membrane bilayers. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1237(2), 183-188.
- Mazzola, C., Medalie, J., Scherma, M., Panlilio, L. V., Solinas, M., Tanda, G., . . . Yasar, S. (2009). Fatty acid amide hydrolase (FAAH) inhibition enhances memory acquisition through activation of PPAR- nuclear receptors. *Learning & Memory*, 16(5), 332-337.

- McAllister, F., Henry, A., Kreindler, J. L., Dubin, P. J., Ulrich, L., Steele, C., . . . Kolls, J. K. (2005). Role of IL-17A, IL-17F, and the IL-17 Receptor in Regulating Growth-Related Oncogene- and Granulocyte Colony-Stimulating Factor in Bronchial Epithelium: Implications for Airway Inflammation in Cystic Fibrosis. *The Journal of Immunology*, 175(1), 404-412.
- McAllister, S. D., Christian, R. T., Horowitz, M. P., Garcia, A., & Desprez, P. (2007). Cannabidiol as a novel inhibitor of Id-1 gene expression in aggressive breast cancer cells. *Molecular Cancer Therapeutics*, 6(11), 2921-2927.
- McAllister, S. D., Murase, R., Christian, R. T., Lau, D., Zielinski, A. J., Allison, J., . . . Desprez, P. (2012). Erratum to: Pathways mediating the effects of cannabidiol on the reduction of breast cancer cell proliferation, invasion, and metastasis. *Breast Cancer Research and Treatment*, 133(1), 401-404.
- Mechoulam, R., & Hanuš, L. (2002). Cannabidiol: An overview of some chemical and pharmacological aspects. Part I: Chemical aspects. *Chemistry and Physics of Lipids*, 121(1-2), 35-43.
- Michalowska, M., Winiarczyk, S., Adaszek, L., Lopuszynski, W., Gradzki, Z., Salmons, B., & Gunzburg, W. (2014). Phase I/II Clinical Trial of Encapsulated, Cytochrome P450 Expressing Cells as Local Activators of Cyclophosphamide to Treat Spontaneous Canine Tumours. *PLOS One*, 9(7), 1-9.
- Minn, A. J., Gupta, G. P., Siegel, P. M., Bos, P. D., Shu, W., Giri, D. D., . . . Massagué, J. (2005). Genes that mediate breast cancer metastasis to lung. *Nature*, 436(7050), 518-524.

- Nakajima, H., Otani, C., & Niimura, T. (1992). Decarboxylation of Gallate by Cell-Free Extracts of *Streptococcus faecalis* and *Klebsiella pneumoniae* Isolated from Rat Feces. *Food Hygiene and Safety Science (Shokuhin Eiseigaku Zasshi)*, 33(4).
- National Cancer Institute. (2017). Cancer Statistics. Retrieved from <https://www.cancer.gov/about-cancer/understanding/statistics>
- Nilsson, M., Hansson, S., Carlsson, A., & Carlsson, M. (2007). Differential effects of the N-methyl-d-aspartate receptor antagonist MK-801 on different stages of object recognition memory in mice. *Neuroscience*, 149(1), 123-130.
- Ozaita, A., Puighermanal, E., & Maldonado, R. (2007). Regulation of PI3K/Akt/GSK-3 pathway by cannabinoids in the brain. *Journal of Neurochemistry*, 102(4), 1105-1114.
- Olea-Herrero, N., Vara, D., Malagarie-Cazenave, S., & Díaz-Laviada, I. (2009). Inhibition of human tumour prostate PC-3 cell growth by cannabinoids R(-)-Methanandamide and JWH-015: Involvement of CB2. *British Journal of Cancer*, 101(6), 940-950.
- Pacher, P., Bátkai, S., & Kunos, G. (2006). The endocannabinoid system as an emerging target of pharmacotherapy. *Pharmacological Reviews*, 58(3), 389-462.
- Pertwee, R. G. (2008a). CB1 and CB2 receptor pharmacology. *Cannabinoids and the Brain*, 91-99.
- Pertwee, R. G. (2008b). The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: Δ^9 -tetrahydrocannabinol, cannabidiol and Δ^9 -tetrahydrocannabivarin. *British Journal of Pharmacology*, 153(2), 199-215.

- Petrocellis, L. D., Ligresti, A., Moriello, A. S., Iappelli, M., Verde, R., Stott, C. G., . . . Marzo, V. D. (2012). Non-THC cannabinoids inhibit prostate carcinoma growth *in vitro* and *in vivo*: Pro-apoptotic effects and underlying mechanisms. *British Journal of Pharmacology*, 168(1), 79-102.
- PharmaCyte. (2017). Live Cell Encapsulation Technology. Retrieved from <http://pharmacYTE.com/live-cell-encapsulation/>
- PharmaCyte. (2017). Cancer Treatment Technologies. Retrieved from <http://pharmacYTE.com/live-cell-encapsulation/cancer/>
- Philip-Ephraim, E. E., Eyong, K. I., Williams, U. E., & Ephraim, R. P. (2012). The role of radiotherapy and chemotherapy in the treatment of primary adult high grade gliomas: Assessment of patients for these treatment approaches and the common immediate side effects. *ISRN Oncology*, 2012, 1-6.
- Pagotto, U., Marsicano, G., Cota, D., Lutz, B., & Pasquali, R. (2006). The emerging role of the endocannabinoid system in endocrine regulation and energy balance. *Endocrine Reviews*, 27(1), 73-100.
- Ramer, R., Merkord, J., Rohde, H., & Hinz, B. (2012). Cannabidiol inhibits cancer cell invasion via upregulation of tissue inhibitor of matrix metalloproteinases-1. *Biochemical Pharmacology*, 79(7), 955-966.
- Reiner, A. M., & Hegeman, G. D. (1971). Metabolism of benzoic acid by bacteria. Accumulation of (-)-3,5-cyclohexadiene-1,2-diol-1-carboxylic acid by a mutant strain of *Alcaligenes eutrophus*. *Biochemistry*, 10(13), 2530-2536.
- Robson, P. (2001). Therapeutic aspects of *Cannabis* and cannabinoids. *The British Journal of Psychiatry*, 178, 107-115.

- Romano, B., Borrelli, F., Pagano, E., Cascio, M. G., Pertwee, R. G., & Izzo, A. A. (2013). Inhibition of colon carcinogenesis by a standardized *Cannabis sativa* extract with high content of cannabidiol. *Phytomedicine*, 21(5), 631-639.
- Sanchez, C., Galve-Roperh, I., Canova, C., Brachet, P., & Guzmán, M. (1998) Delta9-tetrahydrocannabinol induces apoptosis in C6 glioma cells. *FEBS Lett* 436:6–10.
- Salazar, M., Carracedo, A., Salanueva, Í J., Hernández-Tiedra, S., Lorente, M., Egia, A., . . . Velasco, G. (2009). Cannabinoid action induces autophagy-mediated cell death through stimulation of ER stress in human glioma cells. *Journal of Clinical Investigation*, 119(5), 1359-1372.
- Santha, R., Rao, N. A., & Vaidyanathan, C. (1996). Identification of the active-site peptide of 2,3-dihydroxybenzoic acid decarboxylase from *Aspergillus oryzae*. *Biochimica et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology*, 1293(2), 191-200.
- Sharma, M., Hudson, J. B., Adomat, H., Guns, E., & Cox, M. E. (2014). *In vitro* anticancer activity of plant-derived cannabidiol on prostate cancer cell lines. *Pharmacology & Pharmacy*, 05(08), 806-820.
- Scholtzova, H., Wadghiri, Y. Z., Douadi, M., Sigurdsson, E. M., Li, Y., Quartermain, D., . . . Wisniewski, T. (2008). Memantine leads to behavioral improvement and amyloid reduction in Alzheimers-disease-model transgenic mice shown as by micromagnetic resonance imaging. *Journal of Neuroscience Research*, 86(12), 2784-2791.
- Shrivastava, A., Kuzontkoski, P. M., Groopman, J. E., & Prasad, A. (2011). Cannabidiol induces programmed cell death in breast cancer cells by coordinating the cross-

- talk between apoptosis and autophagy. *Molecular Cancer Therapeutics*, 10(7), 1161-1172.
- Snini, S., Tadrist, S., Laffitte, J., Jamin, E., Oswald, I., & Puel, O. (2013). The gene PatG involved in the biosynthesis pathway of patulin, a food-borne mycotoxin, encodes a 6-methylsalicylic acid decarboxylase. *International Journal of Food Microbiology*, 171, 77-83
- Subba, P., Moore, K., & Towers, G. (1967). O-pyrocatechuic acid carboxy-lyase from *Aspergillus niger*. *Archives of Biochemistry and Biophysics*, 122(2), 466-473.
- Thakur, G. A., Duclos, R. I., & Makriyannis, A. (2005). Natural cannabinoids: Templates for drug discovery. *Life Sciences*, 78(5), 454-466.
- Taylor, B., & Ribbons, D. W. (1983). Bacterial Decarboxylation of o-Phthalic Acids. *Applied and Environmental Microbiology*, 46(6), 1276-1281.
- Turu, G., & Hunyady, L. (2010). Signal transduction of the CB1 cannabinoid receptor. *Journal of Molecular Endocrinology*, 44(2), 75-85.
- Xu, X., Liu, Y., Huang, S., Liu, G., Xie, C., Zhou, J., . . . Miao, X. (2006). Overexpression of cannabinoid receptors CB1 and CB2 correlates with improved prognosis of patients with hepatocellular carcinoma. *Cancer Genetics and Cytogenetics*, 171(1), 31-38.

APPENDIX A

CONDITIONS FOR INDIVIDUAL CELL ASSAYS
FOR DECARBOXYLASE ACTIVITY

Table 7: Conditions for Individual Cell Assays for Decarboxylase Activity

Species	Intact or lysed	Induced	Length of Induction	Wet Cell Mass (mg/mL)	Concentration of CBDA (mg/mL)*
<i>Enterobacter (Klebsiella) aerogenes</i>	intact	None	N/A	72.8	10.0
	lysed	None	N/A	72.8	10.0
<i>Pseudomonas putida</i>	intact	2,4-dihydroxybenzoate	1 day	40.4	10.0
	lysed	2,4-dihydroxybenzoate	3 days	40.4	10.0
	lysed	30 mM 2,4-dihydroxybenzoate	1 day	133	22.5
	lysed	30 mM 2,4-dihydroxybenzoate	3 days	69.5	20.0
	lysed	70 mM 2,4-dihydroxybenzoate	1 day	78	20.0
	lysed	10 mM salicylic acid	3 days	68.4	20.0
	lysed	10 mM salicylic acid	3 days	23.6	13.0
	lysed	10 mM salicylic acid	3 days	Not provided	11.0
	lysed	10 mM salicylic acid	3 days	65.5	16.5
	lysed	1 mM salicylic acid	3 days	54.2	2.0
	lysed	1 mM salicylic acid	3 days	58.5	2.0
	lysed	1 mM salicylic acid	3 days	22.3	2.0
<i>Klebsiella pneumonia</i>	intact	30 mM 2,4-dihydroxybenzoate	3 days	162	22.5
	intact	30 mM 2,4-dihydroxybenzoate	3 days	62.4	23.0
	lysed	30 mM 2,4-dihydroxybenzoate	3 days	74.4	23.0
	lysed	None	N/A	84	12.7
	lysed	CBDA	1 day	55.2	12.7
	lysed	30 mM 2,4-dihydroxybenzoate		42.4	15.0
<i>Bacillus Subtilis</i>	lysed	10 mM salicylic acid	2 days	308.8	23.0
	lysed	40 mM salicylic acid	2 days	46	23.0
	lysed	None	N/A	158	12.7
	lysed	CBDA	1 day	206.4	12.7
	lysed	None	N/A	78.1	10.0
	lysed	4 mM salicylic acid	3 days	52.2	10.0
<i>Aspergillus clavatus</i>	lysed	4 mM salicylic acid	3 days	Not provided	10.0
	lysed	4 mM salicylic acid	3 days	Not provided	10.0
<i>Aspergillus niger</i>	lysed	4 mM salicylic acid	3 days	Not provided	2.0
<i>Penicillium chrysogenum</i>	lysed	4 mM salicylic acid	3 days	Not provided	2.0
	lysed	4 mM salicylic acid	3 days	Not provided	13.5
	lysed	4 mM salicylic acid	3 days	Not provided	11.0

*Stock solution in absolute ethanol added to incubation in total of 100 μ L.

APPENDIX B

CHROMATOGRAMS OF FRACTIONS 2-6 COLLECTED
FROM THE PREPARATORY HPLC METHOD

Additional Chromatograms Using the Analytical HPLC Method




Figure 43: Chromatogram of the residue recovered representing peak 2 in the preparatory chromatogram. The sample was resuspended in 1.0 mL absolute ethanol and analyzed using the analytical HPLC method.




Figure 44: Chromatogram of the residue recovered representing peak 3 in the preparatory chromatogram. The sample was resuspended in 1.0 mL absolute ethanol and analyzed using the analytical HPLC method.




Figure 45: Chromatogram of the residue recovered representing peak 4 in the preparatory chromatogram. The sample was resuspended in 1.0 mL absolute ethanol and analyzed using the analytical HPLC method.




Figure 46: Chromatogram of the residue recovered representing peak 5 in the preparatory chromatogram. The sample was resuspended in 1.0 mL absolute ethanol and analyzed using the analytical HPLC method.




Figure 47: Chromatogram of the residue recovered representing peak 6 in the preparatory chromatogram. The sample was resuspended in 1.0 mL absolute ethanol and analyzed using the analytical HPLC method.

APPENDIX C

REPRESENTATIVE HPLC CHROMATOGRAMS FOR
DECARBOXYLASE ACTIVITY IN
VARIOUS CELL LINES

C1. Representative HPLC Chromatograms for the Analysis of *Bacillus subtilis*

The following chromatograms were generated using an isocratic method of 30% HPLC-grade water (solvent A) and 70% HPLC-grade methanol (solvent B).




Figure 48: Chromatogram after 0 hours of incubation of CBDA with lysed *B. subtilis* induced with 40 mM salicylic acid.




Figure 49: Chromatogram after 1 hour of incubation of CBDA with lysed *B. subtilis* induced with 40 mM salicylic acid.




Figure 50: Chromatogram after 2 hours of incubation of CBDA with lysed *B. subtilis* induced with 40 mM salicylic acid.




Figure 51: Chromatogram after 3 hours of incubation of CBDA with lysed *B. subtilis* induced with 40 mM salicylic acid.




Figure 52: Chromatogram after 4 hours of incubation of CBDA with lysed *B. subtilis* induced with 40 mM salicylic acid.




Figure 53: Chromatogram after 0 hours of incubation of CBDA with lysed *B. subtilis* induced with an unknown concentration of CBDA.




Figure 54: Chromatogram after 0.5 hour of incubation of CBDA with lysed *B. subtilis* induced with an unknown concentration of CBDA.




Figure 55: Chromatogram after 1 hour of incubation of CBDA with lysed *B. subtilis* induced with an unknown concentration of CBDA.




Figure 56: Chromatogram after 2 hours of incubation of CBDA with lysed *B. subtilis* induced with an unknown concentration of CBDA.

C2. Representative HPLC Chromatograms for the Analysis of *Enterobacter (Klebsiella) aerogenes*.




Figure 57: Chromatogram after 0 hours of incubation of CBDA with intact *E. aerogenes*.




Figure 58: Chromatogram after 0.5 hour of incubation of CBDA with intact *E. aerogenes*.




Figure 59: Chromatogram after 1 hour of incubation of CBDA with intact *E. aerogenes*.




Figure 60: Chromatogram after 2 hours of incubation of CBDA with intact *E. aerogenes*.




Figure 61: Chromatogram after 3 hours of incubation of CBDA with intact *E. aerogenes*.




Figure 62: Chromatogram after 0 hours of incubation of CBDA with lysed *E. aerogenes*.




Figure 63: Chromatogram after 0.5 hour of incubation of CBDA with lysed *E. aerogenes*.




Figure 64: Chromatogram after 1 hour incubation of CBDA with lysed *E. aerogenes*.




Figure 65: Chromatogram after 2 hours of incubation of CBDA with lysed *E. aerogenes*




Figure 66: Chromatogram after 3 hours of incubation of CBDA with lysed *E. aerogenes*

C3. Representative HPLC Chromatograms for the Analysis of *Klebsiella pneumoniae*

The following chromatograms were generated using an isocratic method of 25% HPLC-grade water (solvent A) and 75% HPLC-grade methanol (solvent B).




Figure 67: CBDA standard for *K. pneumoniae* assay.




Figure 68: Chromatogram after 0 hour of incubation of CBDA with intact *K. pneumoniae* induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.




Figure 69: Chromatogram after 0.5 hour of incubation of CBDA with intact *K. pneumonia* induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.




Figure 70: Chromatogram after 1 hour of incubation of CBDA with intact *K. pneumonia* induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.




Figure 71: Chromatogram after 2 hours of incubation of CBDA with intact *K. pneumonia* induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.




Figure 72: Chromatogram after 3 hours of incubation of CBDA with intact *K. pneumonia* induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.




Figure 73: Chromatogram after 4 hours of incubation of CBDA with intact *K. pneumonia* induced with 30 mM 2,4-dihydroxybenzoic acid for 3 days.

C4. Representative HPLC Chromatograms for the Analysis of *Aspergillus clavatus*

The following chromatograms were generated using an isocratic method of 25% HPLC-grade water (solvent A) and 75% HPLC-grade methanol (solvent B).




Figure 74: Chromatogram after 0 hour of incubation of CBDA with lysed *A. clavatus* Induced with 4 mM salicylic acid.




Figure 75: Chromatogram after 1 hour of incubation of CBDA with lysed *A. clavatus* induced with 4 mM salicylic acid.




Figure 76: Chromatogram after 2 hours of incubation of CBDA with lysed *A. clavatus* induced with 4 mM salicylic acid.




Figure 77: Chromatogram after 3 hours of incubation of CBDA with lysed *A. clavatus* induced with 4 mM salicylic acid.




Figure 78: Chromatogram after 4 hours of incubation of CBDA with lysed *A. clavatus* Induced with 4 mM salicylic acid.

C5. Representative HPLC Chromatograms for the Analysis of *Aspergillus niger*

The following chromatograms were generated using an isocratic method of 25% HPLC-grade water (solvent A) and 75% HPLC-grade methanol (solvent B).




Figure 79: Chromatogram after 0 hour of incubation of CBDA with lysed *A. niger* induced with 4 mM salicylic acid.




Figure 80: Chromatogram after 1 hour of incubation of CBDA with lysed *A. niger* induced with 4 mM salicylic acid.




Figure 81: Chromatogram after 2 hours incubation of CBDA with lysed *A. niger* induced with 4 mM salicylic acid.




Figure 82: Chromatogram after 3 hours of incubation of CBDA with lysed *A. niger* induced with 4 mM salicylic acid.




Figure 83: Chromatogram after 4 hours of incubation of CBDA with lysed *A. niger* induced with 4 mM salicylic acid.

C6. Representative HPLC Chromatograms for the Analysis of *Penicillium chrysogenum*

The following chromatograms were generated using an isocratic method of 25% HPLC-grade water (solvent A) and 75% HPLC-grade methanol (solvent B).




Figure 84: Chromatogram after 0 hours of incubation of CBDA with lysed *P. chrysogenum* induced with 4 mM salicylic acid.




Figure 85: Chromatogram after 1 hour of incubation of CBDA with lysed *P. chrysogenum* induced with 4 mM salicylic acid.




Figure 86: Chromatogram after 2 hours of incubation of CBDA with lysed *P. chrysogenum* induced with 4 mM salicylic acid.




Figure 87: Chromatogram after 3 hours of incubation of CBDA with lysed *P. chrysogenum* induced with 4 mM salicylic acid.




Figure 88: Chromatogram after 4 hours of incubation of CBDA with lysed *P. chrysogenum* induced with 4 mM salicylic acid.


C7. Representative HPLC Chromatograms for the Analysis of *Pseudomonas putida*

Figure 89: Chromatogram after 0 hour of incubation of CBDA with intact *P. putida*. Induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.




Figure 90: Chromatogram after 0.5 hour of incubation of CBDA with intact *P. putida*. Induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.




Figure 91: Chromatogram after 1 hour of incubation of CBDA with intact *P. putida*. Induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.




Figure 92: Chromatogram after 2 hours of incubation of CBDA with intact *P. putida*. Induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.




Figure 93: Chromatogram after 3 hours of incubation of CBDA with intact *P. putida*. Induced 1 day with 30 mM 2,4-dihydroxybenzoic acid.




Figure 94: Chromatogram after 0 hour of incubation of CBDA with lysed *P. putida*. Induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.




Figure 95: Chromatogram after 0.5 hour of incubation of CBDA with lysed *P. putida*. Induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.




Figure 96: Chromatogram after 1 hour of incubation of CBDA with lysed *P. putida*. Induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.




Figure 97: Chromatogram after 2 hours of incubation of CBDA with lysed *P. putida*. Induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.




Figure 98: Chromatogram after 3 hours of incubation of CBDA with lysed *P. putida* induced 3 days with 30 mM 2,4-dihydroxybenzoic acid.

The following chromatograms were generated according to the analytical procedure outlined in Instrumentation of the Methodology section.




Figure 99: Chromatogram after 0 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 100: Chromatogram after 1 hour of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 101: Chromatogram after 2 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 102: Chromatogram after 3 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 103: Chromatogram after 4 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 104: Chromatogram after 5 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 105: Chromatogram after 6 hours of incubation of CBDA with lysed *P. putida* I induced with 10 mM salicylic acid.




Figure 106: Chromatogram after 0 hour of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 107: Chromatogram after 1 hour of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 108: Chromatogram after 2 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 109: Chromatogram after 3 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 110: Chromatogram after 4 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 111: Chromatogram after 0 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 112: Chromatogram after 0.5 hour of incubation of CBDA with lysed *P. putida*. Induced 3 days with 10 mM salicylic acid.




Figure 113: Chromatogram after 1 hour of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 114: Chromatogram after 2 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 115: Chromatogram after 3 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.




Figure 116: Chromatogram after 4 hours of incubation of CBDA with lysed *P. putida* induced with 10 mM salicylic acid.

APPENDIX D

RIGHTS AND PERMISSIONS FOR
USE OF FIGURES

7/17/2018

RightsLink Printable License

JOHN WILEY AND SONS LICENSE TERMS AND CONDITIONS

Jul 17, 2018

This Agreement between Ms. Brittney Morgan ("You") and John Wiley and Sons ("John Wiley and Sons") consists of your license details and the terms and conditions provided by John Wiley and Sons and Copyright Clearance Center.

License Number	4391421292353
License date	Jul 17, 2018
Licensed Content Publisher	John Wiley and Sons
Licensed Content Publication	FEBS Journal
Licensed Content Title	Biosynthesis of cannabinoids
Licensed Content Author	Monika Fellermeier, Wolfgang Eisenreich, Adelbert Bacher, et al
Licensed Content Date	Dec 15, 2003
Licensed Content Volume	268
Licensed Content Issue	6
Licensed Content Pages	9
Type of use	Dissertation/Thesis
Requestor type	University/Academic
Format	Electronic
Portion	Figure/table
Number of figures/tables	1
Original Wiley figure/table number(s)	1
Will you be translating?	No
Title of your thesis / dissertation	AN INVESTIGATION OF ENZYMES CAPABLE OF ACTIVATING CANNABINOID PRODRUGS
Expected completion date	Jul 2018
Expected size (number of pages)	100
Requestor Location	Ms. Brittney Morgan 1411 17th st GREELEY, CO 80631 United States Attn: Ms. Brittney Morgan
Publisher Tax ID	EU826007151
Total	0.00 USD
Terms and Conditions	

TERMS AND CONDITIONS

This copyrighted material is owned by or exclusively licensed to John Wiley & Sons, Inc. or one of its group companies (each a "Wiley Company") or handled on behalf of a society with which a Wiley Company has exclusive publishing rights in relation to a particular work

7/17/2018

RightsLink Printable License

(collectively "WILEY"). By clicking "accept" in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the billing and payment terms and conditions established by the Copyright Clearance Center Inc., ("CCC's Billing and Payment terms and conditions"), at the time that you opened your RightsLink account (these are available at any time at <http://myaccount.copyright.com>).

Terms and Conditions

- The materials you have requested permission to reproduce or reuse (the "Wiley Materials") are protected by copyright.
- You are hereby granted a personal, non-exclusive, non-sub licensable (on a stand-alone basis), non-transferable, worldwide, limited license to reproduce the Wiley Materials for the purpose specified in the licensing process. This license, **and any CONTENT (PDF or image file) purchased as part of your order**, is for a one-time use only and limited to any maximum distribution number specified in the license. The first instance of republication or reuse granted by this license must be completed within two years of the date of the grant of this license (although copies prepared before the end date may be distributed thereafter). The Wiley Materials shall not be used in any other manner or for any other purpose, beyond what is granted in the license. Permission is granted subject to an appropriate acknowledgement given to the author, title of the material/book/journal and the publisher. You shall also duplicate the copyright notice that appears in the Wiley publication in your use of the Wiley Material. Permission is also granted on the understanding that nowhere in the text is a previously published source acknowledged for all or part of this Wiley Material. Any third party content is expressly excluded from this permission.
- With respect to the Wiley Materials, all rights are reserved. Except as expressly granted by the terms of the license, no part of the Wiley Materials may be copied, modified, adapted (except for minor reformatting required by the new Publication), translated, reproduced, transferred or distributed, in any form or by any means, and no derivative works may be made based on the Wiley Materials without the prior permission of the respective copyright owner. **For STM Signatory Publishers clearing permission under the terms of the STM Permissions Guidelines only, the terms of the license are extended to include subsequent editions and for editions in other languages, provided such editions are for the work as a whole in situ and does not involve the separate exploitation of the permitted figures or extracts**, You may not alter, remove or suppress in any manner any copyright, trademark or other notices displayed by the Wiley Materials. You may not license, rent, sell, loan, lease, pledge, offer as security, transfer or assign the Wiley Materials on a stand-alone basis, or any of the rights granted to you hereunder to any other person.
- The Wiley Materials and all of the intellectual property rights therein shall at all times remain the exclusive property of John Wiley & Sons Inc, the Wiley Companies, or their respective licensors, and your interest therein is only that of having possession of and the right to reproduce the Wiley Materials pursuant to Section 2 herein during the continuance of this Agreement. You agree that you own no right, title or interest in or to the Wiley Materials or any of the intellectual property rights therein. You shall have no rights hereunder other than the license as provided for above in Section 2. No right, license or interest to any trademark, trade name, service mark or other branding ("Marks") of WILEY or its licensors is granted hereunder, and you agree that you shall not assert any such right, license or interest with respect thereto

7/17/2018

RightsLink Printable License

- NEITHER WILEY NOR ITS LICENSORS MAKES ANY WARRANTY OR REPRESENTATION OF ANY KIND TO YOU OR ANY THIRD PARTY, EXPRESS, IMPLIED OR STATUTORY, WITH RESPECT TO THE MATERIALS OR THE ACCURACY OF ANY INFORMATION CONTAINED IN THE MATERIALS, INCLUDING, WITHOUT LIMITATION, ANY IMPLIED WARRANTY OF MERCHANTABILITY, ACCURACY, SATISFACTORY QUALITY, FITNESS FOR A PARTICULAR PURPOSE, USABILITY, INTEGRATION OR NON-INFRINGEMENT AND ALL SUCH WARRANTIES ARE HEREBY EXCLUDED BY WILEY AND ITS LICENSORS AND WAIVED BY YOU.
- WILEY shall have the right to terminate this Agreement immediately upon breach of this Agreement by you.
- You shall indemnify, defend and hold harmless WILEY, its Licensors and their respective directors, officers, agents and employees, from and against any actual or threatened claims, demands, causes of action or proceedings arising from any breach of this Agreement by you.
- IN NO EVENT SHALL WILEY OR ITS LICENSORS BE LIABLE TO YOU OR ANY OTHER PARTY OR ANY OTHER PERSON OR ENTITY FOR ANY SPECIAL, CONSEQUENTIAL, INCIDENTAL, INDIRECT, EXEMPLARY OR PUNITIVE DAMAGES, HOWEVER CAUSED, ARISING OUT OF OR IN CONNECTION WITH THE DOWNLOADING, PROVISIONING, VIEWING OR USE OF THE MATERIALS REGARDLESS OF THE FORM OF ACTION, WHETHER FOR BREACH OF CONTRACT, BREACH OF WARRANTY, TORT, NEGLIGENCE, INFRINGEMENT OR OTHERWISE (INCLUDING, WITHOUT LIMITATION, DAMAGES BASED ON LOSS OF PROFITS, DATA, FILES, USE, BUSINESS OPPORTUNITY OR CLAIMS OF THIRD PARTIES), AND WHETHER OR NOT THE PARTY HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES. THIS LIMITATION SHALL APPLY NOTWITHSTANDING ANY FAILURE OF ESSENTIAL PURPOSE OF ANY LIMITED REMEDY PROVIDED HEREIN.
- Should any provision of this Agreement be held by a court of competent jurisdiction to be illegal, invalid, or unenforceable, that provision shall be deemed amended to achieve as nearly as possible the same economic effect as the original provision, and the legality, validity and enforceability of the remaining provisions of this Agreement shall not be affected or impaired thereby.
- The failure of either party to enforce any term or condition of this Agreement shall not constitute a waiver of either party's right to enforce each and every term and condition of this Agreement. No breach under this agreement shall be deemed waived or excused by either party unless such waiver or consent is in writing signed by the party granting such waiver or consent. The waiver by or consent of a party to a breach of any provision of this Agreement shall not operate or be construed as a waiver of or consent to any other or subsequent breach by such other party.
- This Agreement may not be assigned (including by operation of law or otherwise) by you without WILEY's prior written consent.
- Any fee required for this permission shall be non-refundable after thirty (30) days from receipt by the CCC.

7/17/2018

RightsLink Printable License

- These terms and conditions together with CCC's Billing and Payment terms and conditions (which are incorporated herein) form the entire agreement between you and WILEY concerning this licensing transaction and (in the absence of fraud) supersedes all prior agreements and representations of the parties, oral or written. This Agreement may not be amended except in writing signed by both parties. This Agreement shall be binding upon and inure to the benefit of the parties' successors, legal representatives, and authorized assigns.
- In the event of any conflict between your obligations established by these terms and conditions and those established by CCC's Billing and Payment terms and conditions, these terms and conditions shall prevail.
- WILEY expressly reserves all rights not specifically granted in the combination of (i) the license details provided by you and accepted in the course of this licensing transaction, (ii) these terms and conditions and (iii) CCC's Billing and Payment terms and conditions.
- This Agreement will be void if the Type of Use, Format, Circulation, or Requestor Type was misrepresented during the licensing process.
- This Agreement shall be governed by and construed in accordance with the laws of the State of New York, USA, without regards to such state's conflict of law rules. Any legal action, suit or proceeding arising out of or relating to these Terms and Conditions or the breach thereof shall be instituted in a court of competent jurisdiction in New York County in the State of New York in the United States of America and each party hereby consents and submits to the personal jurisdiction of such court, waives any objection to venue in such court and consents to service of process by registered or certified mail, return receipt requested, at the last known address of such party.

WILEY OPEN ACCESS TERMS AND CONDITIONS

Wiley Publishes Open Access Articles in fully Open Access Journals and in Subscription journals offering Online Open. Although most of the fully Open Access journals publish open access articles under the terms of the Creative Commons Attribution (CC BY) License only, the subscription journals and a few of the Open Access Journals offer a choice of Creative Commons Licenses. The license type is clearly identified on the article.

The Creative Commons Attribution License

The Creative Commons Attribution License (CC-BY) allows users to copy, distribute and transmit an article, adapt the article and make commercial use of the article. The CC-BY license permits commercial and non-

Creative Commons Attribution Non-Commercial License

The Creative Commons Attribution Non-Commercial (CC-BY-NC) License permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes. (see below)

Creative Commons Attribution-Non-Commercial-NoDerivs License

The Creative Commons Attribution Non-Commercial-NoDerivs License (CC-BY-NC-ND) permits use, distribution and reproduction in any medium, provided the original work is properly cited, is not used for commercial purposes and no modifications or adaptations are made. (see below)

Use by commercial "for-profit" organizations

Use of Wiley Open Access articles for commercial, promotional, or marketing purposes requires further explicit permission from Wiley and will be subject to a fee.

7/17/2018

RightsLink Printable License

Further details can be found on Wiley Online Library
<http://olabout.wiley.com/WileyCDA/Section/id-410895.html>

Other Terms and Conditions:**v1.10 Last updated September 2015**

Questions? customer care@copyright.com or +1-855-239-3415 (toll free in the US) or
+1-978-646-2777.

7/17/2018

RightsLink Printable License

ELSEVIER LICENSE TERMS AND CONDITIONS

Jul 17, 2018

This Agreement between Ms. Brittney Morgan ("You") and Elsevier ("Elsevier") consists of your license details and the terms and conditions provided by Elsevier and Copyright Clearance Center.

License Number	4391430758400
License date	Jul 17, 2018
Licensed Content Publisher	Elsevier
Licensed Content Publication	Life Sciences
Licensed Content Title	Natural cannabinoids: Templates for drug discovery
Licensed Content Author	Ganesh A. Thakur, Richard I. Duclos, Alexandros Makriyannis
Licensed Content Date	Dec 22, 2005
Licensed Content Volume	78
Licensed Content Issue	5
Licensed Content Pages	13
Start Page	454
End Page	466
Type of Use	reuse in a thesis/dissertation
Intended publisher of new work	other
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
Format	electronic
Are you the author of this Elsevier article?	No
Will you be translating?	No
Original figure numbers	1
Title of your thesis/dissertation	AN INVESTIGATION OF ENZYMES CAPABLE OF ACTIVATING CANNABINOID PRODRUGS
Expected completion date	Jul 2018
Estimated size (number of pages)	100
Requestor Location	Ms. Brittney Morgan 1411 17th st GREELEY, CO 80631 United States Attn: Ms. Brittney Morgan
Publisher Tax ID	98-0397604
Total	0.00 USD

Terms and Conditions

INTRODUCTION

1. The publisher for this copyrighted material is Elsevier. By clicking "accept" in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the Billing and Payment terms and conditions established by Copyright Clearance Center, Inc. ("CCC"), at the time that you opened your Rightslink account and that are available at any time at <http://myaccount.copyright.com>).

GENERAL TERMS

2. Elsevier hereby grants you permission to reproduce the aforementioned material subject to the terms and conditions indicated.

3. Acknowledgement: If any part of the material to be used (for example, figures) has appeared in our publication with credit or acknowledgement to another source, permission must also be sought from that source. If such permission is not obtained then that material may not be included in your publication/copies. Suitable acknowledgement to the source must be made, either as a footnote or in a reference list at the end of your publication, as follows:

"Reprinted from Publication title, Vol /edition number, Author(s), Title of article / title of chapter, Pages No., Copyright (Year), with permission from Elsevier [OR APPLICABLE SOCIETY COPYRIGHT OWNER]." Also Lancet special credit - "Reprinted from The Lancet, Vol. number, Author(s), Title of article, Pages No., Copyright (Year), with permission from Elsevier."

4. Reproduction of this material is confined to the purpose and/or media for which permission is hereby given.

5. Altering/Modifying Material: Not Permitted. However figures and illustrations may be altered/adapted minimally to serve your work. Any other abbreviations, additions, deletions and/or any other alterations shall be made only with prior written authorization of Elsevier Ltd. (Please contact Elsevier at permissions@elsevier.com). No modifications can be made to any Lancet figures/tables and they must be reproduced in full.

6. If the permission fee for the requested use of our material is waived in this instance, please be advised that your future requests for Elsevier materials may attract a fee.

7. Reservation of Rights: Publisher reserves all rights not specifically granted in the combination of (i) the license details provided by you and accepted in the course of this licensing transaction, (ii) these terms and conditions and (iii) CCC's Billing and Payment terms and conditions.

8. License Contingent Upon Payment: While you may exercise the rights licensed immediately upon issuance of the license at the end of the licensing process for the transaction, provided that you have disclosed complete and accurate details of your proposed use, no license is finally effective unless and until full payment is received from you (either by publisher or by CCC) as provided in CCC's Billing and Payment terms and conditions. If full payment is not received on a timely basis, then any license preliminarily granted shall be deemed automatically revoked and shall be void as if never granted. Further, in the event that you breach any of these terms and conditions or any of CCC's Billing and Payment terms and conditions, the license is automatically revoked and shall be void as if never granted. Use of materials as described in a revoked license, as well as any use of the materials beyond the scope of an unrevoked license, may constitute copyright infringement and publisher reserves the right to take any and all action to protect its copyright in the materials.

9. Warranties: Publisher makes no representations or warranties with respect to the licensed material.

10. Indemnity: You hereby indemnify and agree to hold harmless publisher and CCC, and their respective officers, directors, employees and agents, from and against any and all claims arising out of your use of the licensed material other than as specifically authorized pursuant to this license.

11. **No Transfer of License:** This license is personal to you and may not be sublicensed, assigned, or transferred by you to any other person without publisher's written permission.
12. **No Amendment Except in Writing:** This license may not be amended except in a writing signed by both parties (or, in the case of publisher, by CCC on publisher's behalf).
13. **Objection to Contrary Terms:** Publisher hereby objects to any terms contained in any purchase order, acknowledgment, check endorsement or other writing prepared by you, which terms are inconsistent with these terms and conditions or CCC's Billing and Payment terms and conditions. These terms and conditions, together with CCC's Billing and Payment terms and conditions (which are incorporated herein), comprise the entire agreement between you and publisher (and CCC) concerning this licensing transaction. In the event of any conflict between your obligations established by these terms and conditions and those established by CCC's Billing and Payment terms and conditions, these terms and conditions shall control.
14. **Revocation:** Elsevier or Copyright Clearance Center may deny the permissions described in this License at their sole discretion, for any reason or no reason, with a full refund payable to you. Notice of such denial will be made using the contact information provided by you. Failure to receive such notice will not alter or invalidate the denial. In no event will Elsevier or Copyright Clearance Center be responsible or liable for any costs, expenses or damage incurred by you as a result of a denial of your permission request, other than a refund of the amount(s) paid by you to Elsevier and/or Copyright Clearance Center for denied permissions.

LIMITED LICENSE

The following terms and conditions apply only to specific license types:

15. **Translation:** This permission is granted for non-exclusive world **English** rights only unless your license was granted for translation rights. If you licensed translation rights you may only translate this content into the languages you requested. A professional translator must perform all translations and reproduce the content word for word preserving the integrity of the article.
16. **Posting licensed content on any Website:** The following terms and conditions apply as follows: Licensing material from an Elsevier journal: All content posted to the web site must maintain the copyright information line on the bottom of each image; A hyper-text must be included to the Homepage of the journal from which you are licensing at <http://www.sciencedirect.com/science/journal/xxxxx> or the Elsevier homepage for books at <http://www.elsevier.com>; Central Storage: This license does not include permission for a scanned version of the material to be stored in a central repository such as that provided by Heron/XanEdu.
- Licensing material from an Elsevier book: A hyper-text link must be included to the Elsevier homepage at <http://www.elsevier.com>. All content posted to the web site must maintain the copyright information line on the bottom of each image.

Posting licensed content on Electronic reserve: In addition to the above the following clauses are applicable: The web site must be password-protected and made available only to bona fide students registered on a relevant course. This permission is granted for 1 year only. You may obtain a new license for future website posting.

17. **For journal authors:** the following clauses are applicable in addition to the above:

Preprints:

A preprint is an author's own write-up of research results and analysis, it has not been peer-reviewed, nor has it had any other value added to it by a publisher (such as formatting, copyright, technical enhancement etc.).

Authors can share their preprints anywhere at any time. Preprints should not be added to or enhanced in any way in order to appear more like, or to substitute for, the final versions of articles however authors can update their preprints on arXiv or RePEc with their Accepted Author Manuscript (see below).

If accepted for publication, we encourage authors to link from the preprint to their formal publication via its DOI. Millions of researchers have access to the formal publications on ScienceDirect, and so links will help users to find, access, cite and use the best available version. Please note that Cell Press, The Lancet and some society-owned have different preprint policies. Information on these policies is available on the journal homepage.

Accepted Author Manuscripts: An accepted author manuscript is the manuscript of an article that has been accepted for publication and which typically includes author-incorporated changes suggested during submission, peer review and editor-author communications.

Authors can share their accepted author manuscript:

- immediately
 - via their non-commercial person homepage or blog
 - by updating a preprint in arXiv or RePEc with the accepted manuscript
 - via their research institute or institutional repository for internal institutional uses or as part of an invitation-only research collaboration work-group
 - directly by providing copies to their students or to research collaborators for their personal use
 - for private scholarly sharing as part of an invitation-only work group on commercial sites with which Elsevier has an agreement
- After the embargo period
 - via non-commercial hosting platforms such as their institutional repository
 - via commercial sites with which Elsevier has an agreement

In all cases accepted manuscripts should:

- link to the formal publication via its DOI
- bear a CC-BY-NC-ND license - this is easy to do
- if aggregated with other manuscripts, for example in a repository or other site, be shared in alignment with our hosting policy not be added to or enhanced in any way to appear more like, or to substitute for, the published journal article.

Published journal article (JPA): A published journal article (PJA) is the definitive final record of published research that appears or will appear in the journal and embodies all value-adding publishing activities including peer review co-ordination, copy-editing, formatting, (if relevant) pagination and online enrichment.

Policies for sharing publishing journal articles differ for subscription and gold open access articles:

Subscription Articles: If you are an author, please share a link to your article rather than the full-text. Millions of researchers have access to the formal publications on ScienceDirect, and so links will help your users to find, access, cite, and use the best available version.

Theses and dissertations which contain embedded PJAs as part of the formal submission can be posted publicly by the awarding institution with DOI links back to the formal publications on ScienceDirect.

If you are affiliated with a library that subscribes to ScienceDirect you have additional private sharing rights for others' research accessed under that agreement. This includes use for classroom teaching and internal training at the institution (including use in course packs and courseware programs), and inclusion of the article for grant funding purposes.

Gold Open Access Articles: May be shared according to the author-selected end-user license and should contain a [CrossMark logo](#), the end user license, and a DOI link to the formal publication on ScienceDirect.

Please refer to Elsevier's [posting policy](#) for further information.

18. **For book authors** the following clauses are applicable in addition to the above:

Authors are permitted to place a brief summary of their work online only. You are not

allowed to download and post the published electronic version of your chapter, nor may you scan the printed edition to create an electronic version. **Posting to a repository:** Authors are permitted to post a summary of their chapter only in their institution's repository.

19. Thesis/Dissertation: If your license is for use in a thesis/dissertation your thesis may be submitted to your institution in either print or electronic form. Should your thesis be published commercially, please reapply for permission. These requirements include permission for the Library and Archives of Canada to supply single copies, on demand, of the complete thesis and include permission for Proquest/UMI to supply single copies, on demand, of the complete thesis. Should your thesis be published commercially, please reapply for permission. Theses and dissertations which contain embedded PJAs as part of the formal submission can be posted publicly by the awarding institution with DOI links back to the formal publications on ScienceDirect.

Elsevier Open Access Terms and Conditions

You can publish open access with Elsevier in hundreds of open access journals or in nearly 2000 established subscription journals that support open access publishing. Permitted third party re-use of these open access articles is defined by the author's choice of Creative Commons user license. See our [open access license policy](#) for more information.

Terms & Conditions applicable to all Open Access articles published with Elsevier:

Any reuse of the article must not represent the author as endorsing the adaptation of the article nor should the article be modified in such a way as to damage the author's honour or reputation. If any changes have been made, such changes must be clearly indicated. The author(s) must be appropriately credited and we ask that you include the end user license and a DOI link to the formal publication on ScienceDirect.

If any part of the material to be used (for example, figures) has appeared in our publication with credit or acknowledgement to another source it is the responsibility of the user to ensure their reuse complies with the terms and conditions determined by the rights holder.

Additional Terms & Conditions applicable to each Creative Commons user license:

CC BY: The CC-BY license allows users to copy, to create extracts, abstracts and new works from the Article, to alter and revise the Article and to make commercial use of the Article (including reuse and/or resale of the Article by commercial entities), provided the user gives appropriate credit (with a link to the formal publication through the relevant DOI), provides a link to the license, indicates if changes were made and the licensor is not represented as endorsing the use made of the work. The full details of the license are available at <http://creativecommons.org/licenses/by/4.0>.

CC BY NC SA: The CC BY-NC-SA license allows users to copy, to create extracts, abstracts and new works from the Article, to alter and revise the Article, provided this is not done for commercial purposes, and that the user gives appropriate credit (with a link to the formal publication through the relevant DOI), provides a link to the license, indicates if changes were made and the licensor is not represented as endorsing the use made of the work. Further, any new works must be made available on the same conditions. The full details of the license are available at <http://creativecommons.org/licenses/by-nc-sa/4.0>.

CC BY NC ND: The CC BY-NC-ND license allows users to copy and distribute the Article, provided this is not done for commercial purposes and further does not permit distribution of the Article if it is changed or edited in any way, and provided the user gives appropriate credit (with a link to the formal publication through the relevant DOI), provides a link to the license, and that the licensor is not represented as endorsing the use made of the work. The full details of the license are available at <http://creativecommons.org/licenses/by-nc-nd/4.0>. Any commercial reuse of Open Access articles published with a CC BY NC SA or CC BY NC ND license requires permission from Elsevier and will be subject to a fee.

Commercial reuse includes:

- Associating advertising with the full text of the Article
- Charging fees for document delivery or access

11/1/2016

RIGHTSLINK Private License

- Article aggregation
- Systematic distribution via e-mail lists or share buttons

Posting or linking by commercial companies for use by customers of those companies.

20. Other Conditions:

v1.9

Questions? customercare@copyright.com or +1-855-239-3415 (toll free in the US) or +1-978-646-2777.

7/17/2018

RightsLink Printable License

**ELSEVIER LICENSE
TERMS AND CONDITIONS**

Jul :

This Agreement between Ms. Brittney Morgan ("You") and Elsevier ("Elsevier") con
your license details and the terms and conditions provided by Elsevier and Copyright
Clearance Center.

License Number	4391441485271
License date	Jul 17, 2018
Licensed Content Publisher	Elsevier
Licensed Content Publication	Autoimmunity Reviews
Licensed Content Title	Cannabinoids and autoimmune diseases: A systematic review
Licensed Content Author	Valeria Katchan, Paula David, Yehuda Shoenfeld
Licensed Content Date	Jun 1, 2016
Licensed Content Volume	15
Licensed Content Issue	6
Licensed Content Pages	16
Start Page	513
End Page	528
Type of Use	reuse in a thesis/dissertation
Intended publisher of new	other

[Print This Page](#)

7/31/2018

RightsLink Printable License

**AMERICAN ASSOCIATION FOR CANCER RESEARCH LICENSE
TERMS AND CONDITIONS**

Jul 31, 2018

This Agreement between Ms. Brittney Morgan ("You") and American Association for Cancer Research ("American Association for Cancer Research") consists of your license details and the terms and conditions provided by American Association for Cancer Research and Copyright Clearance Center.

License Number	4399381314160
License date	Jul 31, 2018
Licensed Content Publisher	American Association for Cancer Research
Licensed Content Publication	Molecular Cancer Therapeutics
Licensed Content Title	Cannabidiol Induces programmed cell death in breast cancer cells by coordinating the crosstalk between apoptosis and autophagy
Licensed Content Author	Ashutosh Shrivastava, Paula M. Kuzontkoski, Jerome Groopman, Anil Prasad
Licensed Content Date	May 12, 2011
Type of Use	Thesis/Dissertation
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
Will you be translating?	no
Circulation	100
Territory of distribution	North America
Title of your thesis / dissertation	AN INVESTIGATION OF ENZYMES CAPABLE OF ACTIVATING CANNABINOID PRODRUGS
Expected completion date	Jul 2018
Estimated size (number of pages)	100
Requestor Location	Ms. Brittney Morgan 1411 17th st GREELEY, CO 80631 United States Attn: Ms. Brittney Morgan
Billing Type	Invoice
Billing Address	Ms. Brittney Morgan 1411 17th st GREELEY, CO 80631 United States Attn: Ms. Brittney Morgan
Total	0.00 USD

<https://s100.copyright.com/AppDispatchServlet>

1/3

7/31/2018

RightsLink Printable License

Terms and Conditions

American Association for Cancer Research (AACR) Terms and Conditions**INTRODUCTION**

The Publisher for this copyright material is the American Association for Cancer Research (AACR). By clicking "accept" in connection with completing this licensing transaction, you agree to the following terms and conditions applying to this transaction. You also agree to the Billing and Payment terms and conditions established by Copyright Clearance Center (CCC) at the time you opened your Rightslink account.

LIMITED LICENSE

The AACR grants exclusively to you, the User, for onetime, non-exclusive use of this material for the purpose stated in your request and used only with a maximum distribution equal to the number you identified in the permission process. Any form of republication must be completed within one year although copies made before then may be distributed thereafter and any electronic posting is limited to a period of one year. Reproduction of this material is confined to the purpose and/or media for which permission is granted. Altering or modifying this material is not permitted. However, figures and illustrations may be minimally altered or modified to serve the new work.

GEOGRAPHIC SCOPE

Licenses may be exercised as noted in the permission process

RESERVATION OF RIGHTS

The AACR reserves all rights not specifically granted in the combination of 1) the license details provided by you and accepted in the course of this licensing transaction, 2) these terms and conditions, and 3) CCC's Billing and Payment terms and conditions.

DISCLAIMER

You may obtain permission via Rightslink to use material owned by AACR. When you are requesting permission to reuse a portion for an AACR publication, it is your responsibility to examine each portion of content as published to determine whether a credit to, or copyright notice of a third party owner is published next to the item. You must obtain permission from the third party to use any material which has been reprinted with permission from the said third party. If you have not obtained permission from the third party, AACR disclaims any responsibility for the use you make of items owned by them.

LICENSE CONTINGENT ON PAYMENT

While you may exercise the rights licensed immediately upon issuance of the license at the end of the licensing process for the transaction, provided that you have disclosed complete and accurate details of your proposed use, no license is finally effective unless and until full payment is received from you, either by the publisher or by the CCC, as provided in CCC's Billing and Payment terms and conditions. If full payment is not received on a timely basis, then any license preliminarily granted shall be deemed automatically revoked and shall be void as if never granted. Further, in the event that you breach any of these terms and conditions, or any of the CCC's Billing and Payment terms and conditions, the license is automatically revoked and shall be void as if never granted. Use of materials as described in a revoked license, as well as any use of the materials beyond the scope of an unrevoked license, may constitute copyright infringement and the publisher reserves the right to take any and all action to protect its copyright in the materials.

COPYRIGHT NOTICE

You must include the following credit line in connection with your reproduction of the licensed material: "Reprinted (or adapted) from Publication Title, Copyright Year, Volume/Issue, Page Range, Author, Title of Article, with permission from AACR".

TRANSLATION

This permission is granted for non-exclusive world English rights only.

WARRANTIES

Publisher makes no representations or warranties with respect to the licensed material.

7/31/2018

RightsLink Printable License

INDEMNIFICATION

You hereby indemnify and agree to hold harmless the publisher and CCC, and their respective officers, directors, employees and agents, from and against any and all claims arising out of your use of the licensed material other than as specifically authorized pursuant to this license.

REVOCATION

The AACR reserves the right to revoke a license for any reason, including but not limited to advertising and promotional uses of AACR content, third party usage and incorrect figure source attribution.

NO TRANSFER OF LICENSE

This license is personal to you and may not be sublicensed, assigned, or transferred by you to any other person without publisher's written permission.

NO AMENDMENT EXCEPT IN WRITING

This license may not be amended except in a writing signed by both parties (or, in the case of publisher, by CCC on publisher's behalf).

OBJECTION TO CONTRARY TERMS

Publishers hereby objects to any terms contained in any purchase order, acknowledgement, check endorsement or other writing prepared by you, which terms are inconsistent with these terms and conditions or CCC's Billing and Payment terms and conditions. These terms and conditions together with CCC's Billing and Payment terms and conditions (which are incorporated herein) comprise the entire agreement between you and publisher (and CCC) concerning this licensing transaction. In the event of any conflict between your obligations established by these terms and conditions, and those established by CCC's Billing and Payment terms and conditions, these terms and conditions shall control.

THESIS/DISSERTATION TERMS

If your request is to reuse an article authored by you and published by the AACR in your dissertation/thesis, your thesis may be submitted to your institution in either in print or electronic form. Should your thesis be published commercially, please reapply.

ELECTRONIC RESERVE

If this license is made in connection with a course, and the Licensed Material or any portion thereof is to be posted to a website, the website is to be password protected and made available only to the students registered for the relevant course. The permission is granted for the duration of the course. All content posted to the website must maintain the copyright information notice.

JURISDICTION

This license transaction shall be governed by and construed in accordance with the laws of Pennsylvania. You hereby agree to submit to the jurisdiction of the federal and state courts located in Pennsylvania for purposes of resolving any disputes that may arise in connection with this licensing transaction.

Other Terms and Conditions:

v1.0

Questions? customercare@copyright.com or +1-855-239-3415 (toll free in the US) or +1-978-646-2777.